

EXTRACELLULAR MATRIX: TISSUE-SPECIFIC REGULATOR OF CELL PROLIFERATION

AYLIN RIZKI and MINA J. BISSELL

Life Sciences Division, Lawrence Berkeley National Laboratory,
Berkeley, CA 94720

INTRODUCTION

There are two broad categories of extracellular matrix (ECM) in tissues: interstitial/stromal matrix and basement membrane (BM). The interstitial matrix is the loose material around the epithelial cells that is separated from the cells by a basement membrane in many solid tissues. BM is most commonly found lining epithelial cell layers in tissues such as skin and breast (Fuchs et al., 1997; Ronnov-Jessen et al., 1996). Both the composition and the ultrastructure of BM exhibit tissue specificity, as well as temporal regulation during development (Jones and Jones, 2000; Miosge, 2001; Streuli, 1999; Tsai, 1998). Studies of the ultrastructural composition of basement membranes in vivo suggest that the relative arrangement of various ECM components are not only tissue specific but can also be different in certain parts of the same tissue (Lin and Bissell, 1993; Miosge, 2001). For example, ultrastructurally identical basement membranes, such as those found in the proximal and distal tubules of the kidney, have been shown to have a different molecular arrangement when examined by electron microscopy that allows observation of component orientation in tissue samples (Miosge, 2001).

One manifestation of tissue specificity is observed in the form of gene expression patterns, including expression of genes involved in cell cycle regulation. Establishment of tissue-specific gene expression patterns is not simply a result of which ECM molecules surround the cells in the adult tissue. Developmental processes (both during embryogenesis and postbirth, as is the case for the mammary gland) that produce a differentiated tissue involve sequential and interrelated gene regulatory

events that are temporally regulated and that result in an integrated pattern of gene expression in the differentiated tissue (Bissell et al., 1999; Boudreau and Jones, 1999; Lohi et al., 1997; Wiseman and Werb, 2002). Therefore, to fully solve the puzzle of how ECM affects tissue-specific gene expression or cell cycle progression, the information context needs to include the history of the cell and its surrounding ECM (both of which change during development), as well as of the molecular characteristics of the cell-ECM interactions (Bissell and Ram, 1989; Werb and Chin, 1998; Wiseman and Werb, 2002).

Developmental events create a particular imprint in each tissue of both specific ECM molecules and ECM receptors. Thus how a cell behaves in response to its surrounding ECM is dependent not only on the level and composition of the ECM, but also on the cell-surface receptors that recognize and respond to it. The best-studied ECM receptors are the integrin family (Hynes, 1987, 1992; Miranti and Brugge, 2002). However, increasingly other receptors such as syndecans and dystroglycan have been shown to play a role in ECM-mediated signaling (Carey, 1997; Couchman and Woods, 1999; Rapraeger, 2000; Zimmermann and David, 1999). Multiple receptors can recognize a single type of ECM molecule, and a single receptor type may respond to multiple ECM components (Ashkenas et al., 1996; Boudreau and Jones, 1999; Watt, 2002). In addition how a cell responds to ECM is dependent on its growth factor and cytokine context. This is due to the extensive and reciprocal cross-talk between ECM, growth factor, and cytokine receptors (Damsky and Werb, 1992; Danen and Yamada, 2001; Schwartz et al., 1995). Besides cell cycle progression and differentiation, cell-ECM interactions regulate other cellular events such as apoptosis (Boudreau et al., 1995; Howlett et al., 1995). Not surprisingly, disruption of cell-ECM interactions, either by misregulated receptor function or by aberrant ECM composition and arrangement, results in tumorigenesis (Bissell and Radisky, 2001; Radisky et al., 2002; Simpson et al., 1994; Sternlicht et al., 2000; Sternlicht et al., 1999; Talhouk et al., 1992).

TISSUE SPECIFICITY OF ECM AND ITS RECEPTORS

Tissue-specific effects of ECM on cell proliferation are dependent on the molecular composition of the matrix surrounding the cells, as well as the ECM receptor makeup of the particular cell type within a tissue. Here we briefly discuss the function of some of the main ECM component families and their receptors with emphasis on tissue-distribution and tissue-specific diseases associated with these molecules. The ECM components we focus on are collagens, laminins, nidogens, glycosaminoglycans, and proteoglycans; ECM receptors include integrins, dystroglycan, and syndecans. Examples of genetic diseases associated with aberrant ECM or ECM receptor components are listed in Table 9.1.

TABLE 9.1. ECM Component Mutations in Human Disease

| Gene ^a | Disease ^b | Reference ^c |
|------------------------------|--|------------------------------|
| Laminins | | |
| LAMA2 | Merosin-deficient muscular dystrophy | Kuang et al. (1998) |
| LAMA3 | Herlitz type junctional epidermolysis bullosa (skin) | Vidal et al. (1995) |
| LAMB3 | Herlitz type junctional epidermolysis bullosa (skin) | Kon et al. (1998) |
| LAMC3 | Herlitz type junctional epidermolysis bullosa (skin) | Kon et al. (1998) |
| Collagens (fibrillar) | | |
| COL1A1 | Osteogenesis imperfecta (bone, muscle) | Ward et al. (2001) |
| COL1A2 | Osteogenesis imperfecta (bone, muscle) | Trummer et al. (2001) |
| COL1A2 | Ehlers-Danlos syndrome (connective tissue) | Byers et al. (1997) |
| COL1A2 | Marfan syndrome (bone, ocular, cardiovascular) | Dalgleish et al. (1986) |
| COL2A1 | Spondyloepiphyseal dysplasia (bone, retina) | Tiller et al. (1995) |
| COL2A1 | Kniest dysplasia (bone, cartilage) | Spranger et al. (1994) |
| COL2A1 | Achondrogenesis-hypochondrogenesis (bone, cartilage) | Godfrey and Hollister (1988) |
| COL2A1 | Osteoarthritis with mild condrodysplasia (bone, joint) | Ritvaniemi et al. (1995) |
| COL2A1 | Stickler syndrome (joint, hearing, eye, cleft) | Brown et al. (1995) |
| COL2A1 | Wagner syndrome (eye) | Richards et al. (2000) |
| COL3A1 | Ehlers-Danlos syndrome (connective tissue) | Smith et al. (1997) |
| COL3A1 | Familial aortic aneurysms (endothelial) | van Keulen et al. |
| COL6A1 | Bethlem myopathy (muscle) | Jobsis et al. (1996) |
| COL6A2 | Bethlem myopathy (muscle) | Jobsis et al. (1996) |
| COL6A3 | Bethlem myopathy (muscle) | Jobsis et al. (1996) |
| COL10A1 | Metaphyseal chondrodysplasia (bone) | McIntosh et al. (1995) |
| COL10A1 | Spondylomethaphyseal dysplasia (bone) | Ikegawa et al. (1998) |
| COL11A1 | Stickler syndrome (joint, hearing, eye, cleft) | Snead et al. (1996) |
| COL11A1 | Marshall syndrome (hearing, eye, facial skeletal defect) | Meisler et al. (1998) |
| Collagens (BM) | | |
| COL4A3 | Alport syndrome (kidney) | Kashtan (1995) |
| COL4A3 | Benign hematuria (kidney) | Badenas et al. (2002) |
| COL4A4 | Alport syndrome (kidney) | Kashtan (1995) |
| COL4A4 | Benign hematuria (kidney) | Lemmink et al. (1996) |
| COL4A5 | Alport syndrome (kidney) | Kashtan (1995) |
| COL4A6 | Alport syndrome (kidney) | Kashtan (1995) |
| COL8A2 | Fuchs endothelial corneal dystrophy | Biswas et al. (2001) |

^a A gene is listed more than once if it has multiple associated diseases. These genes were selected because mutations in them are associated with genetic diseases.

^b The affected tissues are listed in parentheses.

^c Additional information and references are available at the NIH web site OMIM (Online Mendelian Inheritance In Man), at <http://www.ncbi.nlm.nih.gov:80/entrez/query.fcgi?db=OMIM>.

Collagens

Collagens are the most abundant components of the extracellular matrix as well as of interstitial/stromal matrices. Indeed, collagens are the most abundant proteins in mammals, being a major component of the skin and bone. Most collagens are heterotrimeric molecules of alpha chains (more than 25 of which have been described so far) folded into a coiled-coil triple helix. The triple helix can be homotrimeric, as in the case of type II, type III, type IV, and type VII. Collagens can also be heterotrimeric containing either two or three different alpha chains. For example, collagen type XI is heterotrimeric, containing an alpha 1 (COL11A1), an alpha 2 (COL11A2), and an alpha 3 (COL11A3) chain; collagen type I contains two alpha 1 chains (COL1A1) and one alpha 2 chain (COL1A2). More than 20 different types of collagens have been described based on the structural domains of their alpha chain components. However, within one collagen type, there may be significant variation among molecules because of the large number of alpha chain isoforms that exist. For example, the basement membrane collagen, collagen type IV, can be composed of helix combinations that are compiled from six different gene products (COL4A1 through COL4A6), and their many alternatively spliced variants, in a somewhat tissue-specific manner (Myllyharju and Kivirikko, 2001).

Collagens can be organized into fibrillar structures in connective tissues and in interstitial matrices of soft tissues, or they organize into sheet-like structures in basement membranes. Some nonfibrillar collagens are found associated with fibrils, and sometimes also with the basement membrane and are thought to stabilize interactions between the basement membrane and the interstitial stromal matrix. Fibril-forming collagens include collagen types I, II, III, V, VI, IX, X, and XI (Kadler, 1994, 1995). Fibril-associated collagens are collagen types VII, XII, XV, XVI, XIX, and XXI. Basement membrane collagens are collagen types IV and VIII (Fukai et al., 1994). Functions of collagens XIII, XIV, XXII, XXIII, and XXVI have not been well characterized. Collagen type XVII is an unusual collagen since it is a transmembrane protein (Uitto and Pulkkinen, 1996).

A comprehensive overview of the tissue-specific distribution and functions of this vast number of collagens is not within the scope of this review. However, examples below indicate that certain tissues feature some collagens more prominently, and that diseases associated with each collagen type hints at the tissue specificity of expression (Olsen, 1995; Tryggevason, 1995). Examples of tissue-specific functions and diseases of fibrillar collagen are as follows: Collagen type I is a fibrillar collagen, most commonly found in connective tissues such as bone, cartilage, and skin (Cremer et al., 1998). Mutations in collagen type I have been associated with bone diseases such as osteogenesis imperfecta, Ehlers-Danlos syndrome, and idiopathic osteoporosis, as well as causing a particular type of skin tumor called dermafibrosarcoma protuberans (Kuivaniemi et al., 1997). Collagen type II is a fibrillar collagen found in cartilage and in the vitreous

humor of the eye (Cremer et al., 1998; Kuivaniemi et al., 1997). Accordingly mutations in collagen type II are associated with connective tissue disorders such as achondrogenesis, chondrodysplasia, early onset familial osteoarthritis, SED congenita, Langer-Saldino achondrogenesis, Kniest dysplasia, Stickler syndrome type I, and spondyloepimetaphyseal dysplasia Strudwick type. Collagen type III is also a fibrillar collagen that is associated with flexible connective tissues such as that of the skin, lung, and vasculature, often found associated with collagen type I. Accordingly mutations in this collagen are found associated with blood vessel abnormalities, such as aortic and arterial aneurysms, as well as with connective tissue disorders such as Ehlers-Danlos syndrome (Kuivaniemi et al., 1997). Type V collagen is found in tissues containing type I collagen and appears to regulate the assembly of heterotypic fibers composed of both type I and type V collagen. Collagen VI is a major structural component of muscle microfibrils, and mutations in the genes that code for the collagen VI subunits result in the autosomal dominant disorder, Bethlem myopathy. Type X collagen is short fibrillar collagen that is expressed by chondrocytes during ossification. Mutations in collagen type X result in bone diseases such as Schmid-type metaphyseal chondrodysplasia (SMCD) and Japanese-type spondylometaphyseal dysplasia (SMD) (Kuivaniemi et al., 1997). Collagen type XI is a fibrillar collagen which is a minor constituent of cartilage and mutations in this collagen are associated with type II Stickler syndrome and with Marshall syndrome, both connective tissue disorders (Cremer et al., 1998; Kuivaniemi et al., 1997).

Examples of tissue-specific functions and diseases of basement membrane collagens are as follows: Collagen type IV is found in many basement membranes. There are 6 alpha chains that can give rise to a possible 56 combinations of collagen type IV. Furthermore there are multiple isoforms of many of the subunits, providing a large number of possible collagen type IV combinations for establishment of tissue specificity (Kuhn, 1995; Petitclerc et al., 2000). Mutations in the alpha 1 chain of collagen type IV are associated with type II autosomal Alport syndrome (hereditary glomerulonephropathy) and with familial benign hematuria (thin basement membrane disease). Both diseases affect the kidneys, suggesting that the alpha 3, alpha 4, alpha 5 chains (COL4A3, COL4A4, COL4A5) are prominently featured in the kidney basement membranes (Kashtan, 2000). Type VIII collagen is the main component of the corneal epithelium. This collagen has only two alpha subunits but it can exist either as a homo- or a heterotrimer, as a combination of these two subunits. Mutations in collagen type VIII cause corneal endothelial dystrophy, consistent with the specificity/prominence of its function in the cornea (Meek and Fullwood, 2001).

Laminins

Laminins are a family of heterotrimeric extracellular basement membrane glycoproteins. They are composed of three chains, alpha, beta, and gamma, which form a cruciform structure with three short arms (each

from a different chain) and a long arm (composed of all three chains) (Cheng et al., 1997; Yurchenco et al., 1992). Each laminin subunit has multiple functional domains and is encoded by a distinct gene. Expression of laminin encoding genes is regulated at multiple levels, giving rise to different isoforms. Laminin alpha, beta, and gamma chain isoforms can combine to give rise to different laminins. A total of 11 laminins have been described so far (and they were named in the order of their discovery: laminin-1, laminin-2, etc.); however, a very large number of permutations of the 5 alpha, 4 beta, and 3 gamma subunits and their isoforms is possible (Ekblom et al., 1998).

The tissue-specific distribution, and possible distinct functions of the different alpha, beta, and gamma chains and their isoforms remain largely unknown. Some laminin components have wide tissue distributions. For example, laminin beta 1 is expressed in most tissues that produce a basement membrane, and is one of the three chains constituting laminin 1 (alpha 1/LAMA1, beta 1/LAMB1, gamma 1/LAMC1). This was the first laminin isolated from Engelbreth-Holm-Swarm (EHS) tumor (a basement membrane gel isolated from EHS tumors is widely used in reconstitution experiments in culture and is referred to as a laminin-rich basement membrane, lrBM, in the rest of this chapter) (Friedman et al., 1989; Grant et al., 1985; Kleinman et al., 1982; Li et al., 1987). However, some chains are more prominent in certain tissues, reflecting tissue specificity of BM composition. For example, laminin alpha 2 (LAMA2) is prominently expressed in striated muscle (laminins that contain LAMA2 are called merosins), and the significance of LAMA2 function in muscle is exemplified by the causal relationship between mutations in this gene and congenital merosin-deficient muscular dystrophy (Kuang et al., 1998; Vachon et al., 1996; Wewer and Engvall, 1996). Laminin 5 (alpha 3/LAMA3, beta 3/LAMB3, gamma 2/LAMC2) has a significant role in skin keratinocyte function, as shown by the observation that mutations in any one of its three subunits causes Herlitz type junctional epidermolysis bullosa (Kon et al., 1998; Vidal et al., 1995). Another example of laminin chains that show much restricted tissue distribution is beta 2/LAMB2. It is enriched in the basement membrane of muscles at the neuromuscular junctions, kidney glomerulus and vascular smooth muscle (Hunter et al., 1989; Virtanen et al., 1995).

Nidogens/Entactins

Nidogens are sulfated glycoproteins that are found in basement membranes. Two mammalian nidogens have so far been identified, nidogen 1 and nidogen 2. Their function is to bridge the laminin network with the collagen IV network and provide stability of the basement membranes. Nidogen is essential for both embryonic development and maintenance of proper differentiation of adult tissues. Since there are only two members of the family, nidogen is a less likely candidate for establishment and maintenance of tissue specificity (Dziadek, 1995; Mayer et al., 1998; Timpl et al., 1984). However, distribution of the two members of this family does appear to show some tissue specificity: nidogen 2 is more

prominent in endothelial cells but nidogen 1 is more widely distributed (Schymeinsky et al., 2002).

Proteoglycans and Glycosaminoglycans

Proteoglycans have important structural, as well as signaling functions (Perrimon and Bernfield, 2001). They play a part in providing shape and biomechanical strength to organs and tissues mainly because of the nature of their glycosaminoglycan (GAG) chains. Structural diversity within GAG chains ensures that each protein-GAG interaction is as specific as necessary. A single proteoglycan, even if it carries a single GAG chain, can bind multiple proteins, suggesting the possibility of functional diversity (Delehedde et al., 2001). The core proteins of cell surface proteoglycans may be transmembrane, such as syndecan (an ECM receptor), or GPI-anchored, such as glypican, or matrix proteins, such as perlecan. Perlecan is a heparin sulfate proteoglycan and is a major component of basement membranes (Oldberg et al., 1990; Yanagishita, 1993).

Many growth factors and morphogens (fibroblast growth factors, hepatocyte growth factor/scatter factor, members of the midkine family, and wnts), and matrix proteins (collagen, fibronectin, and laminin) interact with proteoglycans when they signal. The GAG-protein interactions serve to regulate the signal output of growth factor receptor tyrosine kinases and hence cell fate (Park et al., 2000). In addition GAGs coordinate stromal and epithelial development, and they are active participants in mediating cell-cell and cell-matrix interactions (Kresse and Schonherr, 2001; Lander et al., 1996).

An example of a functionally diverse GAG chain (that actually is not attached to protein) is hyaluronic acid/hyaluronan (HA). HA serves a variety of functions, including space filling, lubrication of joints, provision of a matrix through which cells can migrate, and intracellular signaling (Toole, 2001). HA is actively produced during wound healing and tissue repair to provide a framework for growth of blood vessels and fibroblasts (Toole et al., 2002). In addition the interaction of HA with the leukocyte receptor CD44 is important in tissue-specific homing by leukocytes (Turley et al., 2002), and overexpression of HA receptors has been correlated with tumorigenicity and tumor metastasis (Isacke and Yarwood, 2002; Toole, 2002).

Integrins

Integrins are the most extensively studied receptors that transmit ECM signals. The name integrin stems from the observation that integrins are the integrators of extracellular (ECM) and intracellular (cytoskeletal) signals (Boudreau and Jones, 1999; Giancotti and Ruoslahti, 1999; Hynes, 1987). Integrins are heterodimeric receptors of alpha and beta subunits that interact noncovalently to form transmembrane receptors. Currently 17 α and 8 β subunits have been identified. So far, at least 20 heterodimeric receptors have been identified, many of which are tissue specific (Schwartz and Ingber, 1994; Schwartz et al., 1995). Substrates of


integrins are various ECM components including laminins, collagens, fibronectin, and vitronectin. A specific integrin heterodimer can have multiple substrates. For example, $\alpha v \beta 3$ receptor shows strong binding affinity for fibronectin, collagen, tensin-C, thrombospondin, and fibrinogen, as well as vitronectin. A certain type of ECM molecule can interact with more than one kind of integrin heterodimer in a cell context dependent manner (Giancotti and Ruoslahti, 1999). For example, laminin binds $\alpha 3 \beta 1$, $\alpha 6 \beta 1$, and $\alpha 6 \beta 4$ integrins with high affinity. The number of diseases that have been shown to have mutations in integrin subunits is limited so far. This could be partly because such mutations may be lethal. The most prominent defect identified so far is epidermolysis bullosa, a skin disease that is strongly associated with mutations in integrin beta 4 (ITGB4 gene) (Pulkkinen et al., 1998).

In addition to transmitting biochemical signals from ECM proteins, integrins are also involved in sensing and transducing mechanochemical signals (Alenghat and Ingber, 2002; Chen et al., 1997; Ingber, 2002). Integrin signaling has effects on cell adhesion, the cytoskeleton, proliferation, growth, differentiation, and apoptosis pathways. How these pathways are connected is discussed in some detail below, with emphasis on integrin-mediated signaling effects on cell cycle progression.


Non-integrin ECM Receptors

A number of transmembrane cell surface receptor families function in transmitting ECM mediated signals. Among them are syndecans and glypicans. In addition dystroglycan has recently been identified as an ECM signal transmitting receptor. Syndecans are expressed on cells that are dependent on adhesion for proliferation and their distribution is highly cell type and tissue specific. Syndecans are involved in formation

Figure 9.1. Integrin signaling exhibits dynamic reciprocity. (A) ECM-to-nucleus signaling. Integrins are activated and cluster either by crosslinking or occupancy of ECM molecules. Interaction of the cytoplasmic tail of integrins with talin, paxillin, and vinculin or plectin has two main consequences: the actin/keratin cytoskeleton is reorganized, and activation of focal adhesion kinase (FAK) or Shc initiates downstream signaling pathways, such as the Ras-Raf-MEK-ERK (MAPK cascade), the PI-3 kinase, and the JNK pathway. The clustered integrin/ECM and many downstream targets that accumulate at the adhesion site form a focal adhesion complex or a hemidesmosome. Downstream effects of the activation of the MAPK cascade, the PI-3 kinase, and the JNK pathway include transcriptional modulation of a number of genes including those involved in cell cycle regulation. (B) Nucleus-to-ECM signaling. The affinity of membrane-bound integrins for ECM can be altered by change in nuclear functions. Transcriptional changes result in altered expression of cell surface receptors such as integrins, as well as in the composition of ECM via regulation of ECM components and ECM-degrading enzymes (blue). These growth status regulated changes in the nucleus are initiated by the MAPK cascade, or G-protein, PtdIns(4,5)P2 and PKC. MAPKs and PKC signals can also affect the affinity of integrins for their ligands.



A



B




Figure 9.2. Cooperation of integrin and RTK/GFR signaling. (A) ECM activated integrin signaling results in focal adhesion complex formation followed by recruitment of growth factor receptors (GFRs). Such recruitment activates GFRs and their downstream targets such as the MAPK pathway. Alternatively, stability of GFRs can be increased by blocking their ubiquitin-mediated degradation upon attachment to ECM. (B) Sustained activation of the MAPK pathway, as well as activation of RAC-mediated signaling can be dependent on both integrin and GFR signaling.

of focal adhesion complexes, as are integrins (see below), and syndecans can also modulate integrin function (Carey, 1997; Couchman and Woods, 1999; Rapraeger, 2000; Zimmermann and David, 1999).

Glypicans are a distinct family of transmembrane heparan sulfate proteoglycans that contain a core protein anchored to the cell surface via a glycosyl phosphatidylinositol linkage (GPI-linked). Some members of the glypican family of integral membrane proteins are putative cell surface coreceptors for growth factors, ECM proteins, proteases, and antiproteases (Lander et al., 1996), and have been implicated in regulation of cell cycle progression (Delehedde et al., 2001).

Dystroglycan is both an ECM receptor and an organizer of the ECM (Ekblom et al., 1998; Henry et al., 2001). Originally isolated in skeletal muscle, it has now been shown to function in many tissues (Durbeej et al., 1998b). Dystroglycan substrates identified up to now include laminins, and proteoglycans. So far, only one dystroglycan gene has been identified. It has wide tissue distribution and multiple functions (Durbeej and Ekblom, 1997; Durbeej et al., 1998a,b; Muschler et al., 2002).

TISSUE-SPECIFIC EFFECTS OF ECM ON CELL PROLIFERATION IN MOUSE MODELS

Most knockout mouse models of ECM components and ECM receptors are not conditional; therefore the final phenotype observed reflects the

changes throughout development. For this reason, interpretations of such models for tissue specificity of ECM effects on cell cycle progression need to be made with caution. Furthermore ability of another ECM component to substitute for the mutated gene can obscure interpretations. However, using conditional loss-of-function mutations in mice allows observation of tissue-specific developmental defects for certain ECM or receptor components. Tissue-specific conditional knockouts in the adult mouse are the best sources of information for determining cell type specificity of ECM or its receptors in terms of their roles in cell cycle control. Only a few examples of conditional knockouts exist for ECM and its receptors, namely for $\beta 1$ integrin and dystroglycan (Brakebusch et al., 2000; Hirsch et al., 1996; Raghavan et al., 2000). An overview of ECM and ECM receptor components that have been knocked out in mice is shown in Table 9.2.

Examination of homozygous knockouts of ECM components and their receptors shows that loss of a single subunit is usually not embryonic lethal, suggesting that other BM components can substitute for function in most tissues. Nidogen-1 and nidogen-2 knockout mice demonstrate an extreme case of this. Targeted homozygous disruption of neither nidogen-1 nor nidogen-2 results in the disruption of BM. Since nidogen is necessary for the integrity of basement membranes, this suggests that nidogen-1 and nidogen-2 can substitute for each other's function with no or very little tissue specific requirements (Murshed et al., 2000; Schymeinsky et al., 2002). In other ECM knockouts, some tissues and cell types show gross developmental abnormalities, some of which are due to changes in the regulation of cell proliferation by ECM. For example, disruption of the alpha 3 chain of laminin 5 result in abnormalities in the survival of epithelial cells (especially in the skin) as well as extreme blistering of the skin, similar to a skin disease known as junctional epidermolysis bullosa-gravis in humans. LAMA3 was shown to be necessary for the proper formation and stabilization of hemidesmosomes in the epidermis, and the skin abnormalities could be attributed to the hemidesmosome defects (Ryan et al., 1999). A few knockouts do show embryonic lethality, suggesting that the mutated genes are essential in early development and not replaceable by similar components. For example, mice homozygous null for the gamma 1 chain of laminin (LAMC1), beta 1 integrin (ITGB1), dystroglycan (DAG1) show embryonic lethality (Smyth et al., 1999; Stephens et al., 1995; Williamson et al., 1997). Conditional knockouts of integrin beta 1 and dystroglycan have been produced for multiple tissues (Brakebusch et al., 2000; Cohn et al., 2002; Hirsch et al., 1996; Moore et al., 2002; Raghavan et al., 2000). Conditional knockouts are more informative about cell proliferation effects especially when the knockout event occurs either after completion of the development of the tissue or in later stages of development. For example, disrupting beta 1 integrin in skin, using a keratin 14 promoter that is turned on very late in development, results in failure of basement membrane assembly and maintenance, and impairment of epidermal proliferation (Raghavan et al., 2000).

TABLE 9.2. ECM and ECM Receptor Mutations in Mice

| Gene | Homozygous Mutant Phenotype | Reference |
|----------------------|---|--|
| Laminins | | |
| LAMA2 | Murine muscular dystrophy | Xu et al. (1994) |
| LAMA3 | Neonatal lethality | Ryan et al. (1999) |
| LAMA4 | Nervous system developmental defects | Patton et al. (2001) |
| LAMB2 | Aberrant neuromuscular development | Noakes et al. (1995) |
| LAMB3 | Junctional epidermolysis bullosa (skin blistering) | Kuster et al. (1997) |
| LAMC1 | Embryonic lethality | Smyth et al. (1999) |
| Collagens | | |
| COLA1 | Dermal fibrosis, impaired uterine involution | Liu et al. (1995) |
| COLA2 | Glomerulopathy (kidney) | Phillips et al. (2002) |
| COL3A1 | Cardiovascular defects | Liu et al. (1997) |
| COL4A3 | Alport syndrome (kidney) | Cosgrove et al. (1996) |
| COL5A2 | Spinal deformities, skin and eye abnormalities | Andrikopoulos et al. (1995) |
| COL10A1 | Defects in bone development and hematopoiesis | Gress and Jacenko (2000) |
| COL11A1 | Skeletal development defects | Li et al. (1995) |
| COL11A2 | Hearing loss | McGuirt et al. (1999) |
| COL15A1 | Skeletal myopathy, cardiovascular defects | Eklund et al. (2001) |
| Nidogens | | |
| NID1 | No detectable defects | Murshed et al. (2000) |
| NID2 | No detectable defects | Schymeinsky et al. (2002) |
| Proteoglycans | | |
| HSPG2 | Cartilage defects, cardiac and brain BM defects | Costell et al. (1999) |
| AGC | Bone and cartilage phenotypes | Wai et al. (1998) |
| FN1 | Mesoderm, neural tube, vascular development defects | George et al. (1993) |
| Integrins | | |
| ITGA1 | Derived MEFs have adhesion defects in vitro | Gardner et al. (1996) |
| ITGA2 | Kidney and lung development defects | Kreidberg et al. (1996) |
| ITGA4 | Placental and cardiac development defects | Yang et al. (1995) |
| ITGA5 | Embryonic mesodermal defects | Yang et al. (1993) |
| ITGA6 | Epidermolysis bullosa (skin blisters), neonatal death | Georges-Labouesse et al. (1996) |
| ITGA7 | Homozygous null shows muscle and tendon organization defects, including a novel form of muscular dystrophy | Mayer et al. (1997), Miosge et al. (1999) |
| ITGA8 | Homozygous null has defects in kidney morphogenesis | Muller et al. (1997) |
| ITGA9 | Homozygous mutants have fatal bilateral chylothorax (lymphatic system and thoracic duct defects) | Huang et al. (2000) |
| ITGAE | Cutaneous inflammatory disorder | Schon et al. (2000) |
| ITGAM | Neutrophil adhesion, migration, apoptosis defects | Coxon et al. (1996), Ding et al. (1999) |
| ITGAV | Lethality preceded by aberrant vasculogenesis, angiogenesis, and organogenesis | Bader et al. (1998) |
| ITGB1 | Peri-implantation lethality and inner cell mass failure Their ES cells have migration and adhesion defects | Fassler et al. (1995), Stephens et al. (1995) |

TABLE 9.2. *Continued*

| Gene | Homozygous Mutant Phenotype | Reference |
|---------------------|--|---|
| ITGB1 | Conditional null in skin has epidermal proliferation, basement membrane formation, and hair follicle invagination defects | Brakebusch et al. (2000), Raghavan et al. (2000) |
| ITGB1 | Chimeric mice, with null hematopoietic cells display impaired migration of hematopoietic stem cells | Hirsch et al. (1996) |
| ITGB3 | Homozygous mutants of the cytoplasmic domain have impaired platelet function | Law et al. (1999) |
| ITGB3 | Osteoclast abnormalities | McHugh et al. (2000) |
| ITGB3 | Heterozygous mutants have placental defects and reduced survival | Hodivala-Dilke et al. (1999) |
| ITGB4 | Homozygous nulls have defects in hemidesmosome formation, cell adhesion, and cell survival, followed by lethality soon after birth | Dowling et al. (1996), Frei et al. (1999) |
| ITGB4 | Homozygous deletion of the cytoplasmic domain results in cell cycle and adhesion defects, and lethality postbirth | Murgia et al. (1998) |
| ITGB6 | Juvenile baldness and asthma-like syndromes due to infiltration of skin and lung epithelium by macrophages and lymphocytes | Huang et al. (1996) |
| ITGB7 | Defects in gut-associated lymphoid tissue formation possibly due to defective lymphocyte attachment | Wagner et al. (1996) |
| ITGA8 | Homozygous nulls die soon after birth due to kidney defects, with ear neuroepithelial deformities | Littlewood Evans and Muller (2000) |
| Syndecans | | |
| SDC1 | Decreased Wnt-1 induced mammary tumorigenesis | Alexander et al. (2000) |
| SDC3 | Feeding behavior defects due to altered hypothalamic functions (predominantly neural) | Reizes et al. (2001) |
| SDC4 | Homozygous mutants display delayed healing of skin wounds and defective angiogenesis | Echtermeyer et al. (2001) |
| Glypicans | | |
| GPC3 | Defects in limb patterning, skeletal development, and kidney branching morphogenesis, renal cystic dysplasia, ventral wall defects | Cano-Gauci et al. (1999), Grisaru et al. (2001), Paine-Saunders et al. (2000) |
| Dystroglycan | | |
| DAG1 | Early embryonic lethality associated with defects in the extra-embryonic basement membrane | Williamson et al. (1997) |
| DAG1 | Conditional mutants with homozygous deletion in the brain display congenital muscular dystrophy syndromes | Moore et al. (2002) |
| DAG1 | Conditional mutants with homozygous deletion in skeletal muscle display muscle regeneration defects | Cohn et al. (2002) |

Note: The *Homozygous Mutant Phenotype* column shows the phenotype of mice carrying a targeted homozygous null mutation of the indicated gene, unless otherwise indicated to be a conditional mutation in a certain tissue or a heterozygous mutation.




Figure 9.3. ECM-initiated signals regulate G₁ and S phase entry and progression. Sustained activation of Erk via integrin-mediated and/or RTK initiated activation of the MAPK pathway results in translocation of Erk into the nucleus. Erk regulates cyclin D1 expression positively and p21 and p27 negatively. Activation of cyclin D1-cdk4/6 complexes allow G₁ entry. Progression through G₁ is controlled by cyclin E-cdk2. Cyclin E-cdk2 activation requires downregulation of p21 or p27, which is accomplished by Erk-mediated signaling as well. Activation of cyclin E-cdk2 and of cyclin D-cdk4/6 free up the E2F transcription factor by phosphorylating Rb. E2F is needed for transcription of cyclin A. Thus S phase entry and progression that is dependent on cyclin A-cdk2 activation is also regulated by ECM, since downstream targets of cyclin D1 activation and p21 or p27 inactivation include cyclin A. In addition to the MAPK pathway, PI-3K or Rac can also upregulate cyclin D1 transcription through activation of JNK via Akt (a downstream target of PI-3K) or via JNK (a downstream target of Rac).

EFFECTS OF CELL-ECM INTERACTIONS ON CELL CYCLE PROGRESSION IN CULTURED CELLS

Overview

For most cells derived from multicellular organisms, proliferation *ex vivo* requires simulating at least part of the cellular microenvironment within the tissue. Many nonmalignant cells derived from solid, multicellular organs require adhesion to a substratum in order to proliferate. ECM components, such as fibronectin, laminin, and collagen, have been used as thin monolayer coats onto which cells can attach and grow (Ashkenas et al., 1996). Although plastic that is treated to simulate a

charge distribution that allows cellular attachment has been widely used, presence of ECM provides additional biochemical signals that can promote or inhibit cell proliferation (Cambier et al., 2000; Giancotti and Ruoslahti, 1999; Schwartz, 2001). The ECM effect on cell cycle is dependent on cell type, cell-surface receptor composition, as well as the nature of the ECM molecules.

A classic example of positive regulation of growth by ECM, which was later followed by many other examples, is the demonstration that mouse bone marrow cells, when cultured on a complex ECM derived from marrow, exhibit a dramatically increased ability to proliferate, compared to the controls (Campbell et al., 1985). Negative regulation of growth by ECM in nontumorigenic adherent cells is perhaps best demonstrated by use of three-dimensional (3D) cultures of laminin rich reconstituted basement membrane (IrBM). For example, mammary epithelial cells plated in 3D IrBM proliferate for only a limited number of divisions, arrest growth, and form differentiated colonies resembling tissue structures in vivo, in both shape and function (Fig. 9.4) (Petersen et al., 1992), and specific integrins such as $\alpha\beta 8$ exert negative growth control in epithelial cells (Cambier et al., 2000). Loss of dependence on adherence or acquisition of anchorage-independent growth (as measured by growth in soft agar) usually accompanies transformation to malignancy. Interestingly, tumorigenic counterparts of cell types that differentiate within 3D ECM cultures exhibit loss of growth regulation as well as their ability to differentiate in response to ECM-mediated signals in the same assay.

Control of Cell Proliferation by Integrin Signaling

ECM-mediated signals that are transmitted via integrins are generally involved in controlling events in G_1 phase progression. Of the G_1 phase cyclins and CDKs (cyclin D, cyclin E, CDK2 and CDK4 and their inhibitors), integrin signaling has the most significant effect on the induction of cyclin D1 and repression of the CDK inhibitors p21 and p27 (Assoian 1997; Roovers and Assoian 2000) (Fig. 9.3). Signaling through integrins is bi-directional (Bissell et al., 1982; Chen et al., 1994). ECM signals are transmitted through integrins to their downstream effectors that regulate expression and activity of the cell cycle regulators in the nucleus. For the rest of this review we call these events ECM-to-nucleus signaling. Growth signals from the nucleus reciprocally regulate the extracellular levels and binding potential of ECM molecules and integrins by regulating their structure and function. We refer to such events as nucleus-to-ECM signaling. While we have focused on integrins as the main transducers of such bi-directional flow of information, or dynamic reciprocity, between the nucleus and cell surface receptors, this phenomenon is also observed for growth factor, cytokine, and other receptors.

ECM-to-Nucleus Signaling. In response to ECM signaling, integrins and many associated proteins cluster and associate with the cellular cytoskeleton to promote filament assembly or disassembly, followed by an intricate series of signaling events that result in changes in a number




Figure 9.4. Nontumorigenic and tumorigenic mammary epithelial cells differ in their ability to proliferate and differentiate in lrBM. Tritiated thymidine incorporation was measured as a function of time in primary "normal" or nontumorigenic mammary epithelial cell lines (**A**), and in tumorigenic lines (**B**). Initially both normal and tumorigenic cells are able to divide. However, normal cells growth arrest and differentiate (**A**), but tumorigenic cells continue proliferating (**B**) (Reprinted with modifications by permission of Cancer Biology from Weaver et al. 1995). (See color insert.)

of nuclear responses including transcription of cell cycle regulators (Fig. 9.1A). The overview below describes the main components of such signaling events, each of which is a potential point of regulation for cell cycle progression.

Reorganization of actin or keratin filaments into large stress fibers produces signals that feed back into the integrin clusters and causes enhanced integrin clustering and increased binding to the ECM. This region of clustering and signaling is called either a focal adhesion if actin reorganization is involved or a hemidesmosome if the keratin cytoskeleton is involved as in the case of $\alpha 6 \beta 4$ signaling (Jones et al., 1998; Nievers

et al., 1999). Although for the rest of this section we focus on focal adhesion-mediated signaling, similar principles also apply to hemidesmosome-mediated signaling. Integrin clustering can further be affected by lateral association of integrins with other membrane proteins, such as caveolin-1 and urokinase plasminogen activator (uPAR), an ECM degrading enzyme (Wary et al., 1996; Wei et al., 1999). Integrin clustering activates some protein tyrosine kinases, including focal adhesion kinase (FAK), Src-family kinases, Abl, as well as a serine-threonine kinase called integrin-linked kinase (ILK) (Giancotti and Ruoslahti, 1999; Wary et al., 1996, 1999). First, interaction of the cytoplasmic tail of integrins with cytoskeletal proteins such as talin and paxillin results in recruitment of FAK to the nascent focal adhesions (Chen et al., 1995; Lewis and Schwartz, 1995; Miyamoto et al., 1995; Parsons et al., 1994; Schaller et al., 1995). This is followed by autophosphorylation of FAK, creating a binding site for Src homology 2 (SH2) domain proteins (Schaller et al., 1994; Schlaepfer et al., 1994). The SH2 domain kinase then phosphorylates a number of focal adhesion proteins, including cytoskeletal proteins such as paxillin and tensin, docking proteins such as p130^{CAS} that recruits adapter proteins such as Crk and Nck (Vuori et al., 1996). FAK can activate phosphoinositide 3-OH kinase (PI 3-kinase) either through activation of Src or directly (Chen et al., 1996). The interaction of Src and FAK can be reciprocal; that is, Src can phosphorylate FAK at the same tyrosine that is autophosphorylated by FAK. This creates a binding site for the adapter complex containing Grb2 and Ras guanosine 5'-triphosphate exchange factor mSOS (Schlaepfer et al., 1994). These signals are transmitted to the MAPK pathway through activation cascade of Ras, Raf, MEK, and ERK sequentially. The MAPKs are also downstream targets of growth factor receptor tyrosine kinases, providing a link between integrin and growth factor receptor signaling. The downstream effects of MAPK signaling include controlling cyclin D1 expression (needed for cell cycle entry), as well as controlling integrin and ECM molecule expression in a feedback loop (nucleus-to-ECM signaling).

Another feedback loop is observed between mitotic signals and FAK. FAK is further phosphorylated on serine when cells enter mitosis. This results in dissociation of FAK from Src and p130^{CAS} (Yamakita et al., 1999). Loosening of focal contacts may allow the cells to decrease adhesion to the substratum and help them divide and spread out. A parallel pathway of integrin-mediated activation starts with activation of a Src family kinase, Fyn, by some integrins. In this pathway a membrane-bound receptor (e.g., caveolin-1) acts as an adapter linking the integrin alpha subunit to Fyn. When ECM binding activates integrins, Fyn becomes activated and its Src homology 3 (SH3) domain interacts with Shc, which in turn phosphorylates it on tyrosine. This targets Shc for the adapter Grb2-mSOS complex. The SOS complex then transduces signals to the MAPK pathways through Ras, Raf, MEK, and ERK (Schlaepfer et al., 1994), which in turn induce changes in nuclear events that result in enhanced cell proliferation.

Both activation of FAK and Shc (and perhaps other yet unidentified molecules) can contribute to the Ras-ERK MAPK signaling cascade (Howe et al., 2002; Hughes et al., 1997). In some cell types Shc is responsible for the high-level activation of ERK, following cell adhesion to a substratum. FAK most likely has a more significant role in sustaining the activation signal (Pozzi et al., 1998; Wary et al., 1999). Both activation of FAK and Shc can be regulated positively and negatively by tyrosine phosphatases. Some examples include receptor-type protein phosphatase alpha and cytosolic phosphatase SHP-2, which remove the negative regulatory phosphate in Src kinases and therefore amplify both FAK and Shc signaling (Oh et al., 1999; Su et al., 1999). PTP-PEST and PTP-1B are examples of cytoplasmic tyrosine phosphatases that dephosphorylate p130^{CAS}, and this inhibits some of the downstream signals of FAK (Garton et al., 1997; Garton and Tonks, 1999; Liu et al., 1998). The phosphatases themselves can be regulated by signals that are integrin-ECM activated. For example, PTP-PEST is anchored to the endoplasmic reticulum and needs to be recruited to the focal adhesions. This occurs when integrin-mediated adhesion activates a protease called calpain, which in turn cleaves the PTP-PEST extension and anchors it to the endoplasmic reticulum, allowing the phosphatase to localize to focal adhesions (Rock et al., 1997). Since phosphatases can have multiple specificities, they can affect cell proliferation by regulating two related pathways. For example, PTEN (a tumor-suppressor gene-encoded protein) dephosphorylates PI 3-kinase generated inositol lipids (Li et al., 1997; Maehama and Dixon, 1998; Myers et al., 1998; Stambolic et al., 1998). PTEN can also dephosphorylate FAK and Shc and suppress integrin signaling. Inhibition of PI 3-kinase results in downmodulation of Ras Raf Erk (MAPK) signaling (Gu et al., 1998; Tamura et al., 1998) (Fig. 9.1A). This, in turn, can downmodulate active integrin expression. Therefore a phosphatase like PTEN can inhibit focal adhesion formation via multiple mechanisms. Inhibition of adhesion, as well as down-regulation of the PI 3-kinase survival pathway, can cooperate to reduce the cell's ability to proliferate.

Nucleus-to-ECM Signaling. This can manifest itself in multiple forms (Fig. 9.1B). In many instances, proliferation-related signals that alter nuclear functions result in altered functional expression of cell surface receptors or of the ECM components produced by the cell, thereby changing both the ECM composition surrounding the cell and its ability to respond to the changed microenvironment.

For example, the MAPK-dependent differentiation and growth arrest of PC12 cells is accompanied by up-regulation of $\alpha 1\beta 2$ integrin expression, which helps maintain an elongated morphology via its collagen/laminin interactions (Rossino et al., 1990). A similar nucleus-to-ECM signaling event is observed in erythroleukemia cells that are dependent on MAPK for differentiation; MAPK-mediated growth arrest and differentiation in these cells is accompanied by up-regulation of receptor $\alpha \text{IIb}\beta 3$ (Woods et al., 2001). Furthermore expression of some ECM components is dependent on the MAPK-initiated differentiation. Therefore growth arrest related changes in the nucleus result both in

increased ligand concentration for integrin binding and in expression of the integrin receptors (Jones et al., 1999).

An interesting mechanism by which the proliferation status of cells results in nuclear signals that manifest themselves in the form of altered cell-ECM interactions comes from the studies of the Ets transcription factor PEA3 (polyome enhancer activator 3). PEA3 is a direct downstream target of ERK and is essential for the transcriptional activation of many integrins, including α IIb and α v integrins. PEA3 also plays a significant role in the expression of proteases that degrade ECM, such as uPA (urokinase plasminogen activator), collagenases, and stromelysin-1 (Boudreau and Jones, 1999; Crawford and Matrisian, 1996; Wasylyk et al., 1991).

It is intriguing to note that MAPK activation, when it reflects a change in proliferation, can also lead to suppression of high affinity ligand binding of many integrins including β 1, β 3, and α 6, by a mechanism that does not involve transcriptional or translational regulation of integrins or their ligands directly. In a screen for suppressors of integrin activation, H-Ras and its effector kinase Raf-1 were identified as negative regulators of integrin activation (Hughes et al., 1997). H-Ras inhibited the activation of integrins with three distinct alpha and beta subunit cytoplasmic domains. Suppression was independent of both transcription and protein synthesis. Furthermore suppression correlated with activation of the ERK MAP kinase pathway. Therefore the effect of ECM/adhesion on cell proliferation via integrin activation is tightly interconnected with proliferation signals that alter nuclear events. Such nuclear signals regulate integrin composition, affinity of integrins for their ligands, and the composition of the ligands as demonstrated by the transcriptional effects on ECM molecules themselves, as well as on the expression of ECM degrading enzymes.


Cooperation of Integrins and Growth Factor Receptors. Multiple mechanisms exist for crosstalk between integrin-mediated and growth factor/receptor tyrosine kinase (RTK) signaling that result in a cumulative effect on cell proliferation (Fig. 9.3). Following integrin activation by ECM, some RTKs are recruited to the focal adhesion complexes. In various cell types, epidermal growth factor receptor (EGFR), platelet-derived growth factor receptor (PDGFR), and fibroblast growth factor receptor (FGFR) have been found recruited to focal adhesion complexes (Miyamoto et al., 1996; Plopper and Ingber, 1993; Plopper et al., 1995). This recruitment results in phosphorylation and activation of growth factor receptors and their downstream targets (Miyamoto et al., 1996). Cell adhesion-mediated signals can also affect stability of growth factor receptors as exemplified by an increase in the number of PDGF receptors by blocking ubiquitin-mediated degradation of the receptor following adherence to a fibronectin substratum (Baron and Schwartz, 2000). The downstream MAPK cascade of Ras, Raf, MEK, and ERK can also be differentially affected by adhesion-mediated signaling. There are reported instances where activation of Raf and its downstream targets were promoted by adhesion, but Ras activation was not (Lin et al., 1997;

Roovers and Assoian, 2000). Furthermore integrin-mediated adhesion can be necessary for activation of MEK by Raf and maintenance of active ERK requires adhesion (Howe et al., 2002; Renshaw et al., 1999; Roovers and Assoian, 2000; Roovers et al., 1999). Pathways that are sensitive to both integrin and RTK-initiated signals include the Rac pathway. Growth factor induction of Rac is integrin dependent as demonstrated by observations that RTKs can activate Rac in suspended cells, but the activated Rac is not targeted to the plasma membrane and does not interact and activate its downstream targets such as PI 3-kinase and Akt (del Pozo et al., 2000; Khwaja et al., 1997).

It should be noted that the crosstalk between integrins and RTKs is not only dependent on the presence or absence of integrin-mediated adhesion but also on the nature of the substrata. Experiments with mammary epithelial cells grown either as monolayers on 2D plastic substratum or in 3D laminin-rich reconstituted basement membrane (lrBM) suggest that the composition of the substrata has profound consequences for integrin and growth factor receptor crosstalk. Tumorigenic mammary epithelial cells are unable to growth arrest in 3D lrBM but their normal counterparts stop growth and differentiate into *in vivo* like structures called acini (Petersen et al., 1992; Weaver et al., 1995) (Fig. 9.4). However, blocking cell-ECM interactions by inhibiting $\beta 1$ integrin or inactivating EGF receptor signaling in tumorigenic cells in 3D lrBM results in reacquisition of an ability to growth arrest and differentiate (Wang et al., 1998; Weaver et al., 1997) (Fig. 9.5A and B). In 3D lrBM, inhibiting one receptor ($\beta 1$ integrin or EGFR) results in down-regulation of the expression of the other (EGFR or $\beta 1$ integrin), but this is not the case when cells are grown in monolayer 2D cultures (Fig. 9.5C). Conversely, both EGFR (ErbB1) and ErbB2 can induce increased proliferation in MCF-10A cells in monolayer, but in 3D lrBM, once the acini are formed, only ErbB2 results in uncontrolled proliferation that fills the acinar lumen with cells and ErbB1 has no effect (Muthuswamy et al., 2001). Taken together with the effect of the MAPK pathway on nucleus-to-ECM signaling that can determine integrin composition, integrin affinity, and ECM molecule composition (see above), these observations suggest that the role of context should be taken into account when interpreting signaling events.

Cell Cycle Regulatory Targets of Integrin-Mediated Signaling.

Integrin-mediated signaling can induce either cell cycle progression or exit from the cell cycle followed by differentiation. Positive regulation of cell proliferation by integrins has been observed in cell types that are dependent on adhesion for cell cycle entry. The most common and most highly studied role of integrin signaling on promoting cell cycle entry is in promoting progression to G_1 and the G_1/S transition. The G_1 phase is also controlled by growth factor signals; therefore the crosstalk between integrins and growth factors can manifest itself in the form of co-regulation of cell cycle progression. Although overcoming G_2/M arrest can contribute to cell cycle progression, the role of ECM signaling on the G_2/M transition has not been given much attention. One reason for








| | 3D lrBM | | | | 2D plastic | | |
|--------------------------|---|---|---|----------------|---|---|----------------|
| | S1 | fr4-2 | T4-2 "treated" | | T4-2 | T4-2 "treated" | |
| |  |  |  | |  |  | |
| Inhibitor Added: | — | — | β1 Integrin Inhibitor | EGFR Inhibitor | — | β1 Integrin Inhibitor | EGFR Inhibitor |
| β1 Integrin total levels | + | +++ | + | + | +++ | +++ | +++ |
| EGFR total levels | + | ++++ | + | + | ++++ | ++++ | ++++ |
| EGFR* activated | + | ++++ | + | + | ++++ | ++++ | + |

Figure 9.5. Reverted tumorigenic mammary epithelial cells exhibit crosstalk between $\beta 1$ integrin and EGFR in 3D lrBM but not in 2D monolayer cultures. Treatment of tumorigenic human mammary epithelial cells by inhibitory antibodies against $\beta 1$ integrin or EGFR results in reversion to differentiated acinar phenotype as shown by re-establishment of basal $\alpha 6 \beta 4$ integrin expression. (Reprinted with permission of *Journal of Cell Biology* from Weaver et al., 1997) (A) $\alpha 6$ integrin is mislocalized in tumorigenic T4-2 cells (orange). (See color insert.) (B) $\alpha 6$ integrin is relocalized to the basal side after tumorigenic T4-2 cells are reverted to a differentiated, nondividing structure in lrBM by treating with inhibitory $\beta 1$ integrin antibodies (orange). (C) Inhibiting $\beta 1$ integrin in T4-2 cells results in downregulation of the level and activity of EGFR to levels similar to what is observed for non-tumorigenic S1 cells when T4 is grown in 3D lrBM but not in monolayer 2D cultures. Likewise inhibition of EGFR results in $\beta 1$ integrin down-regulation in 3D but not in 2D. (Reprinted with permission of *Cancer Research* from Bissell et al., 1999).

this is that several studies have shown that only G_1 phase is subject to control by the ECM (and growth factors) for the cell types and the substrata studied (Fang et al., 1996; Sherr, 1994; Zhu et al., 1996). Therefore more thorough studies using a larger number of cell types and substrata are needed to determine if ECM signaling under normal growth conditions may control G_2/M transition. In addition, whether ECM plays a role in the G_2/M progression when cells are arrested in response to external or internal stimuli, such as DNA damaging agents, has not yet been determined.

The mammalian cell cycle is controlled by cyclins, cyclin-dependent kinases (cdks) that are activated by cyclins, cdk inhibitors, and cyclin activators. There are many cdks that function at different stages of the cell cycle. Progression through G_1 phase is modulated by the binding of cyclin D family (D1, D2, D3) to cdk4 or its homolog cdk6 (cdk4/6). Cyclin E binding to cdk2 is also involved in G_1 phase progression. Binding of cyclin A to cdk2 is required for progression through S phase. Activated Cdk4/6 and cdk2 are required for phosphorylation of the retinoblastoma protein (Rb), which activates transcription of genes regulated by E2F family of transcription factors. One of the functions of E2Fs is to regulate expression of cyclin A and therefore S phase entry (Weinberg, 1995). This finalizes the known points of cell cycle regulation initiated by adhesion or growth factors. Progression through G_2/M phase of the cell cycle is activated by binding of cyclin B to cdk1 (=cdc2) (Assoian, 1997; Heichman and Roberts, 1994; Sherr, 1994); however, G_2/M transition does not appear to be one of the main points of regulation initiated by ECM or growth factors. Progression through G_1 or G_2/M can be inhibited by either the INK4 family (p15, p16, p18, and p19) or the p21 family (p21cip1, p27kip1, and p53kip2) of cdk inhibitors. The INK4 inhibitors bind and inhibit cdk4/6, resulting in G_1 arrest. The p21 family of cdk inhibitors can bind to, and inhibit, either cdk4/6 or cdk2, preventing progression to G_1 or S phases of the cell cycle.

The effect of ECM-mediated cell cycle control is mainly manifested by regulation of G_1 and S phase progression for both positive and negative regulatory functions (Schwartz, 2001) (Fig. 9.3). In cases where integrin-mediated adhesion exhibits a positive effect on cell proliferation, activation of the Ras-Raf-MEK-ERK cascade by either ECM-mediated signaling or by growth factors results in increased cyclin D1 expression or down-regulation of the cdk inhibitors p21 and p27 (Roovers and Assoian, 2000). At least in some cases, ERK activity by itself is not sufficient for cyclin D1 expression, cells need to be attached to a substratum (Weber et al., 1997). For example, in suspended CCL39 cells, which require adhesion for proliferation, sustained ERK activity is not sufficient for cyclin D1 activation (Le Gall et al., 1998). It has been reported that translocation of ERK from the cytoplasm to the nucleus is also adhesion dependent (Aplin et al., 2001; Assoian and Schwartz, 2001). It is indicated that other downstream targets that are regulated by both integrin and growth factor receptor/RTK signaling, such as PI 3-kinase, play a role in the transcriptional activation and stability of cyclin D1 (Danilkovitch et al., 2000; Gille and Downward, 1999; Takuwa et al.,

1999). Cyclin D1 regulation by integrin signaling can also be manifested at the level of translation, at least in some cell types, in a manner that is dependent on activation of Rac, a signaling molecule that is translocated to the plasma membrane via events mediated by integrins (Huang et al., 1998; Schwartz and Assoian, 2001). Therefore, in addition to the MAPK signaling cascade, PI 3-kinase and Rac-mediated events can control the transcription and translation of cyclin D1.

Integrin signaling can also mediate cell cycle progression by regulating the cell cycle inhibitors p21 and p27. These cdk inhibitors are normally upregulated in early G₁ phase, but their levels drop as cells progress through G₁ to allow exit from G₁ and entry into the S phase. Induction of p21 in adherence-dependent cells is strongly regulated by ERK activity in early G₁ and the down-regulation of p21 in late G₁ is impaired when integrin signaling is inhibited. However, involvement of ERK with p21 appears to be indirect and through the small GTP binding protein Rho. Regulation of Rho activity by cytoskeletal changes, such as induction of actin stress fibers, has been well studied; however, the impact of integrin signaling on Rho remains somewhat controversial. One reason for the controversy is the difficulty of separating the effects of ECM on cell shape (known to induce Rac) from its effects on integrin signaling. A recent report indicates that activation of the downstream effector of integrin signaling, FAK, can negatively regulate Rho activity (Ren et al., 1999, 2000; Schwartz and Assoian, 2001). Effects of Rho-mediated regulation of cdk inhibitors is likely to turn out to be a cell type, cell surface receptor, and ECM-specific effect, as are many of the signaling cascades described above.

Dependence of cell cycle regulation by ECM on cell type, receptor composition, and substratum identity is best demonstrated by the observations that integrins can also regulate cell cycle progression negatively. For example, expression of $\alpha\text{v}\beta 8$ results in increased expression of the cdk inhibitor p21 in a lung carcinoma cell line that has lost $\alpha\text{v}\beta 8$ with subsequent growth arrest when grown on a monolayer of vitronectin, an $\alpha\text{v}\beta 8$ ligand (Cambier et al., 2000). Many lung tumors have lost their $\alpha\text{v}\beta 8$ and introduction of this integrin to tumorigenic lung epithelial cells results in loss of tumorigenicity in nude mice (Cambier et al., 2000).

CONCLUSION

A large number of studies on the effect of ECM on proliferation of cultured cells, data on tissue-specific diseases that are associated with specific ECM signaling components, and knockout mice studies provide clues that control of cell proliferation within a tissue is dependent on the nature of the ECM and its receptors in the tissues. A more comprehensive understanding of contribution of ECM to the tissue-specific regulation of cell cycle progression will require more systematic approaches, including conditional knockouts in different tissues as well as development of culture systems that better mimic physiological tissue environment.

ACKNOWLEDGMENTS

We thank the numerous investigators, including the past and present members of the Bissell laboratory, who contributed to the field of extracellular matrix research and apologize to those colleagues whose work we were unable to cite due to space limitations. The authors' work was supported by funds from the U.S. Department of Energy, Office of Biological and Environmental Research, the National Cancer Institute, and by an Innovator Award from the U.S. Department of Defense Breast Cancer Research Program (to M.J.B.), and by a Postdoctoral Fellowship from the California Breast Cancer Research Program (to A.R.).

REFERENCES

- Alenghat FJ, Ingber DE (2002): Mechanotransduction: All signals point to cytoskeleton, matrix, and integrins. *Sci STKE* 2002:PE6.
- Alexander CM, Reichsman F, Hinkes MT, et al (2000): Syndecan-1 is required for Wnt-1-induced mammary tumorigenesis in mice. *Nat Genet* 25:329-32.
- Andrikopoulos K, Liu X, Keene DR, Jaenisch R, Ramirez F (1995): Targeted mutation in the *col5a2* gene reveals a regulatory role for type V collagen during matrix assembly. *Nat Genet* 9:31-6.
- Aplin AE, Stewart SA, Assoian RK, Juliano RL (2001): Integrin-mediated adhesion regulates ERK nuclear translocation and phosphorylation of Elk-1. *J Cell Biol* 153:273-82.
- Ashkenas J, Muschler J, Bissell MJ (1996): The extracellular matrix in epithelial biology: Shared molecules and common themes in distant phyla. *Dev Biol* 180: 433-44.
- Assoian RK (1997): Control of the G₁ phase cyclin-dependent kinases by mitogenic growth factors and the extracellular matrix. *Cytokine Growth Factor Rev* 8:165-70.
- Assoian RK, Schwartz MA (2001): Coordinate signaling by integrins and receptor tyrosine kinases in the regulation of G₁ phase cell-cycle progression. *Curr Opin Genet Dev* 11:48-53.
- Badenas C, Praga M, Tazon B, et al (2002): Mutations in the COL4A4 and COL4A3 genes cause familial benign hematuria. *J Am Soc Nephrol* 13: 1248-54.
- Bader BL, Rayburn H, Crowley D, Hynes RO (1998): Extensive vasculogenesis, angiogenesis, and organogenesis precede lethality in mice lacking all alpha v integrins. *Cell* 95:507-19.
- Baron V, Schwartz M (2000): Cell adhesion regulates ubiquitin-mediated degradation of the platelet-derived growth factor receptor beta. *J Biol Chem* 275: 39318-23.
- Bissell MJ, Hall HG, Parry G (1982): How does the extracellular matrix direct gene expression? *J Theor Biol* 99:31-68.
- Bissell MJ, Radisky D (2001): Putting tumours in context. *Nat Rev Cancer* 1: 46-54.
- Bissell MJ, Ram TG (1989): Regulation of functional cytodifferentiation and histogenesis in mammary epithelial cells: Role of the extracellular matrix. *Environ Health Perspect* 80:61-70.

- Bissell MJ, Weaver VM, Lelievre SA, Wang F, Petersen OW, Schmeichel KL (1999): Tissue structure, nuclear organization, and gene expression in normal and malignant breast. *Cancer Res* 59:1757s-63s; Discussion 1763s-64s.
- Biswas S, Munier FL, Yardley J, et al (2001): Missense mutations in COL8A2, the gene encoding the alpha2 chain of type VIII collagen, cause two forms of corneal endothelial dystrophy. *Hum Mol Genet* 10:2415-23.
- Boudreau N, Simpson CJ, Werb Z, Bissell MJ (1995): Suppression of ICE and apoptosis in mammary epithelial cells by extracellular matrix. *Science* 267:891-3.
- Boudreau NJ, Jones PL (1999): Extracellular matrix and integrin signalling: the shape of things to come. *Biochem J* 339 (Pt 3):481-8.
- Brakebusch C, Grose R, Quondamatteo F, et al (2000): Skin and hair follicle integrity is crucially dependent on beta 1 integrin expression on keratinocytes. *EMBO J* 19:3990-4003.
- Brown DM, Vandenburg K, Kimura AE, Weingeist TA, Sheffield VC, Stone EM (1995): Novel frameshift mutations in the procollagen 2 gene (COL2A1) associated with Stickler syndrome (hereditary arthro-ophthalmopathy). *Hum Mol Genet* 4:141-2.
- Byers PH, Duvic M, Atkinson M, et al (1997): Ehlers-Danlos syndrome type VIIA and VIIB result from splice-junction mutations or genomic deletions that involve exon 6 in the COL1A1 and COL1A2 genes of type I collagen. *Am J Med Genet* 72:94-105.
- Cambier S, Mu DZ, O'Connell D, et al (2000): A role for the integrin alpha-v-beta-8 in the negative regulation of epithelial cell growth. *Cancer Res* 60:7084-93.
- Campbell A, Wicha MS, Long M (1985): Extracellular matrix promotes the growth and differentiation of murine hematopoietic cells in vitro. *J Clin Invest* 75:2085-90.
- Cano-Gauci DF, Song HH, Yang H, et al (1999): Glypican-3-deficient mice exhibit developmental overgrowth and some of the abnormalities typical of Simpson-Golabi-Beckwith syndrome. *J Cell Biol* 146:255-64.
- Carey DJ (1997): Syndecans: Multifunctional cell-surface co-receptors. *Biochem J* 327 (Pt 1):1-16.
- Chen CS, Mrksich M, Huang S, Whitesides GM, Ingber DE (1997): Geometric control of cell life and death. *Science* 276:1425-8.
- Chen HC, Appeddu PA, Isoda H, Guan JL (1996): Phosphorylation of tyrosine 397 in focal adhesion kinase is required for binding phosphatidylinositol 3-kinase. *J Biol Chem* 271:26329-34.
- Chen HC, Appeddu PA, Parsons JT, Hildebrand JD, Schaller MD, Guan JL (1995): Interaction of focal adhesion kinase with cytoskeletal protein talin. *J Biol Chem* 270:16995-9.
- Chen YP, O'Toole TE, Shipley T, et al (1994): "Inside-out" signal transduction inhibited by isolated integrin cytoplasmic domains. *J Biol Chem* 269:18307-10.
- Cheng YS, Champlaud MF, Burgeson RE, Marinkovich MP, Yurchenco PD (1997): Self-assembly of laminin isoforms. *J Biol Chem* 272:31525-32.
- Cohn RD, Henry MD, Michele DE, et al (2002): Disruption of DAG1 in differentiated skeletal muscle reveals a role for dystroglycan in muscle regeneration. *Cell* 110:639-48.

- Cosgrove D, Meehan DT, Grunkemeyer JA, et al (1996): Collagen COL4A3 knockout: A mouse model for autosomal Alport syndrome. *Genes Dev* 10: 2981-92.
- Costell M, Gustafsson E, Aszodi A, et al (1999): Perlecan maintains the integrity of cartilage and some basement membranes. *J Cell Biol* 147:1109-22.
- Couchman JR, Woods A (1999): Syndecan-4 and integrins: Combinatorial signaling in cell adhesion. *J Cell Sci* 112 (Pt 20):3415-20.
- Coxon A, Rieu P, Barkalow FJ, et al (1996): A novel role for the beta 2 integrin CD11b/CD18 in neutrophil apoptosis: A homeostatic mechanism in inflammation. *Immunity* 5:653-66.
- Crawford HC, Matrisian LM (1996): Mechanisms controlling the transcription of matrix metalloproteinase genes in normal and neoplastic cells. *Enzyme Protein* 49:20-37.
- Cremer MA, Rosloniec EF, Kang AH (1998): The cartilage collagens: a review of their structure, organization, and role in the pathogenesis of experimental arthritis in animals and in human rheumatic disease. *J Mol Med* 76:275-88.
- Dagleish R, Williams G, Hawkins JR (1986): Length polymorphism in the pro alpha 2(I) collagen gene: An alternative explanation in a case of Marfan syndrome. *Hum Genet* 73:91-2.
- Damsky CH, Werb Z (1992): Signal transduction by integrin receptors for extracellular matrix: cooperative processing of extracellular information. *Curr Opin Cell Biol* 4:772-81.
- Danen EH, Yamada KM (2001): Fibronectin, integrins, and growth control. *J Cell Physiol* 189:1-13.
- Danilkovitch A, Donley S, Skeel A, Leonard EJ (2000): Two independent signaling pathways mediate the antiapoptotic action of macrophage-stimulating protein on epithelial cells. *Mol Cell Biol* 20:2218-27.
- del Pozo MA, Price LS, Alderson NB, Ren XD, Schwartz MA (2000): Adhesion to the extracellular matrix regulates the coupling of the small GTPase Rac to its effector PAK. *EMBO J* 19:2008-14.
- Delehedde M, Lyon M, Sergeant N, Rahmoune H, Fernig DG (2001): Proteoglycans: Pericellular and cell surface multireceptors that integrate external stimuli in the mammary gland. *J Mammary Gland Biol Neoplasia* 6:253-73.
- Ding ZM, Babensee JE, Simon SI, et al (1999): Relative contribution of LFA-1 and Mac-1 to neutrophil adhesion and migration. *J Immunol* 163:5029-38.
- Dowling J, Yu QC, Fuchs E (1996): Beta4 integrin is required for hemidesmosome formation, cell adhesion and cell survival. *J Cell Biol* 134:559-72.
- Durbeej M, Ekblom P (1997): Dystroglycan and laminins: Glycoconjugates involved in branching epithelial morphogenesis. *Expr Lung Res* 23:109-18.
- Durbeej M, Henry MD, Campbell KP (1998a): Dystroglycan in development and disease. *Curr Opin Cell Biol* 10:594-601.
- Durbeej M, Henry MD, Ferletta M, Campbell KP, Ekblom P (1998b): Distribution of dystroglycan in normal adult mouse tissues. *J Histochem Cytochem* 46:449-57.
- Dziadek M (1995): Role of laminin-nidogen complexes in basement membrane formation during embryonic development. *Experientia* 51:901-13.
- Echtermeyer F, Streit M, Wilcox-Adelman S, et al (2001): Delayed wound repair and impaired angiogenesis in mice lacking syndecan-4. *J Clin Invest* 107: R9-R14.




Figure 2.1. Subnuclear compartmentalization of nucleic acids and regulatory proteins into specialized domains. See text for full caption.




Figure 2.3. Scaffolding nuclear proteins: A mechanism of specificity in gene regulation. See text for full caption.




Figure 6.2. Gallery of fluorescent micrographs depicting glutaraldehyde-fixed and lysed PtK₁ cells in various stages of mitosis. See text for full caption.




Figure 9.4. Nontumorigenic and tumorigenic mammary epithelial cells differ in their ability to proliferate and differentiate in IrBM. See text for full caption.

A

T4-2



B

T4-2, $\beta 1$ -blocked

Figure 9.5. Reverted tumorigenic mammary epithelial cells exhibit crosstalk between $\beta 1$ integrin and EGFR in 3D IrBM but not in 2D monolayer cultures. See text for full caption.




Figure 10.1. Continuous versus bolus administration of human endostatin to SCID mice bearing human pancreatic cancer that is $p53^{+/+}$. See text for full caption.




Figure 10.4. In mice depleted of the endogenous angiogenesis inhibitor tumstatin, tumors grow 300% to 400% more rapidly than in wild-type mice. See text for full caption.




Figure 10.8. Tumor cells are genetically unstable and contain thousands of genomic alterations. See text for full caption.




Figure 11.1. Developmental stages of the fruit fly *Drosophila melanogaster*. See text for full caption.




Figure 16.4. (B) Two subpathways exist in nucleotide excision repair, global genome repair, and transcription coupled repair. See text for full caption.




Figure 16.8. Schematic representation of the homologous recombination mechanism. See text for full caption.




Figure 17.2. Activation of the Src oncoprotein. See text for full caption.




Figure 17.3. Activation of the Abl oncoprotein. See text for full caption.




Figure 19.6. Structure of wild-type p53 bound to DNA. Protein Data Bank ID: 1TUP (see Web Resources).




Figure 19.7. Prima-1 reactivates mutant p53. **(A)** Restoration of wild-type p53 activity to mutant p53 by Prima-1 in mouse 10(3) cells. Murine (10)3 fibroblasts lacking endogenous p53 were engineered to express only the selectable marker (CMV) or either the human mutant p53-R175H or R273H. Cells were grown under normal culture conditions (control) or treated with Prima-1 (10 μ M) for 48 hours and stained for morphological analysis. Note that cells lacking p53 maintained viability after Prima-1 treatment (upper right panel), whereas cells expressing mutant p53 underwent apoptosis (middle and lower right panels) (unpublished data). **(B)** Restoration of wild-type p53 activity to mutant p53 by Prima-1 in Saos-2 cells. Human osteosarcoma Saos-2 cells lacking endogenous p53 were engineered to express only the selectable marker (CMV) or human mutant p53-R281G. Cells were grown under normal culture conditions (Control) or treated with Prima-1 (75 μ M) for 48 hours and stained for morphological analysis. Note that cells lacking p53 maintained viability after Prima-1 treatment (upper right panel) whereas cells expressing mutant p53 underwent apoptosis (lower right panel) (unpublished data).

- Ekblom M, Falk M, Salmivirta K, Durbeek M, Ekblom P (1998): Laminin isoforms and epithelial development. *An NY Acad Sci* 857:194-211.
- Eklund L, Piuhola J, Komulainen J, et al (2001): Lack of type XV collagen causes a skeletal myopathy and cardiovascular defects in mice. *Proc Natl Acad Sci USA* 98:1194-9.
- Fang F, Orend G, Watanabe N, Hunter T, Ruoslahti E (1996): Dependence of cyclin E-CDK2 kinase activity on cell anchorage. *Science* 271:499-502.
- Fassler R, Pfaff M, Murphy J, et al (1995): Lack of beta 1 integrin gene in embryonic stem cells affects morphology, adhesion, and migration but not integration into the inner cell mass of blastocysts. *J Cell Biol* 128:979-88.
- Frei R, Dowling J, Carenini S, Fuchs E, Martini R (1999): Myelin formation by Schwann cells in the absence of beta4 integrin. *Glia* 27:269-74.
- Friedman SL, Roll FJ, Boyles J, Arenson DM, Bissell DM (1989): Maintenance of differentiated phenotype of cultured rat hepatic lipocytes by basement membrane matrix. *J Biol Chem* 264:10756-62.
- Fuchs E, Dowling J, Segre J, Lo SH, Yu QC (1997): Integrators of epidermal growth and differentiation: Distinct functions for beta 1 and beta 4 integrins. *Curr Opin Genet Dev* 7:672-82.
- Fukai N, Apte SS, Olsen BR (1994): Nonfibrillar collagens. *Methods Enzymol* 245:3-28.
- Gardner H, Kreidberg J, Koteliensky V, Jaenisch R (1996): Deletion of integrin alpha 1 by homologous recombination permits normal murine development but gives rise to a specific deficit in cell adhesion. *Dev Biol* 175:301-13.
- Garton AJ, Burnham MR, Bouton AH, Tonks NK (1997): Association of PTP-PEST with the SH3 domain of p130cas; a novel mechanism of protein tyrosine phosphatase substrate recognition. *Oncogene* 15:877-85.
- Garton AJ, Tonks NK (1999): Regulation of fibroblast motility by the protein tyrosine phosphatase PTP-PEST. *J Biol Chem* 274:3811-8.
- George EL, Georges-Labouesse EN, Patel-King RS, Rayburn H, Hynes RO (1993): Defects in mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin. *Development* 119:1079-91.
- Georges-Labouesse E, Messaddeq N, Yehia G, Cadalbert L, Dierich A, Le Meur M (1996): Absence of integrin alpha 6 leads to epidermolysis bullosa and neonatal death in mice. *Nat Genet* 13:370-3.
- Giancotti FG, Ruoslahti E (1999): Integrin signaling. *Science* 285:1028-32.
- Gille H, Downward J (1999): Multiple ras effector pathways contribute to G(1) cell cycle progression. *J Biol Chem* 274:22033-40.
- Godfrey M, Hollister DW (1988): Type II achondrogenesis-hypochondrogenesis: Identification of abnormal type II collagen. *Am J Hum Genet* 43:904-13.
- Grant DS, Kleinman HK, Leblond CP, Inoue S, Chung AE, Martin GR (1985): The basement-membrane-like matrix of the mouse EHS tumor: II. Immunohistochemical quantitation of six of its components. *Am J Anat* 174:387-98.
- Gress CJ, Jacenko O (2000): Growth plate compressions and altered hematopoiesis in collagen X null mice. *J Cell Biol* 149:983-93.
- Grisaru S, Cano-Gauci D, Tee J, Filmus J, Rosenblum ND (2001): Glypican-3 modulates BMP- and FGF-mediated effects during renal branching morphogenesis. *Dev Biol* 231:31-46.
- Gu J, Tamura M, Yamada KM (1998): Tumor suppressor PTEN inhibits integrin- and growth factor-mediated mitogen-activated protein (MAP) kinase signaling pathways. *J Cell Biol* 143:1375-83.

- Heichman KA, Roberts JM (1994): Rules to replicate by. *Cell* 79:557-62.
- Henry MD, Satz JS, Brakebusch C, et al. (2001): Distinct roles for dystroglycan, beta1 integrin and perlecan in cell surface laminin organization. *J Cell Sci* 114:1137-44.
- Hirsch E, Iglesias A, Potocnik AJ, Hartmann U, Fassler R (1996): Impaired migration but not differentiation of haematopoietic stem cells in the absence of beta1 integrins. *Nature* 380:171-5.
- Hodivala-Dilke KM, McHugh KP, Tsakiris DA, et al. (1999): Beta3-integrin-deficient mice are a model for Glanzmann thrombasthenia showing placental defects and reduced survival. *J Clin Invest* 103:229-38.
- Howe AK, Aplin AE, Juliano RL (2002): Anchorage-dependent ERK signaling—Mechanisms and consequences. *Curr Opin Genet Dev* 12:30-5.
- Howlett AR, Bailey N, Damsky C, Petersen OW, Bissell MJ (1995): Cellular growth and survival are mediated by beta 1 integrins in normal human breast epithelium but not in breast carcinoma. *J Cell Sci* 108 (Pt 5):1945-57.
- Huang S, Chen CS, Ingber DE (1998): Control of cyclin D1, p27(Kip1), and cell cycle progression in human capillary endothelial cells by cell shape and cytoskeletal tension. *Mol Biol Cell* 9:3179-93.
- Huang XZ, Wu JF, Cass D, et al. (1996): Inactivation of the integrin beta 6 subunit gene reveals a role of epithelial integrins in regulating inflammation in the lung and skin. *J Cell Biol* 133:921-8.
- Huang XZ, Wu JF, Ferrando R, et al. (2000): Fatal bilateral chylothorax in mice lacking the integrin alpha9beta1. *Mol Cell Biol* 20:5208-15.
- Hughes PE, Renshaw MW, Pfaff M, et al. (1997): Suppression of integrin activation: A novel function of a Ras/Raf-initiated MAP kinase pathway. *Cell* 88:521-30.
- Hunter DD, Shah V, Merlie JP, Sanes JR (1989): A laminin-like adhesive protein concentrated in the synaptic cleft of the neuromuscular junction. *Nature* 338:229-34.
- Hynes RO (1987): Integrins: A family of cell surface receptors. *Cell* 48:549-54.
- Hynes RO (1992): Integrins: Versatility, modulation, and signaling in cell adhesion. *Cell* 69:11-25.
- Ikegawa S, Nishimura G, Nagai T, Hasegawa T, Ohashi H, Nakamura Y (1998): Mutation of the type X collagen gene (COL10A1) causes spondylometaphyseal dysplasia. *Am J Hum Genet* 63:1659-62.
- Ingber D (2002): Mechanical signaling. *An NY Acad Sci* 961:162-3.
- Isacke CM, Yarwood H (2002): The hyaluronan receptor, CD44. *Int J Biochem Cell Biol* 34:718-21.
- Jobsis GJ, Keizers H, Vreijling JP, et al. (1996): Type VI collagen mutations in Bethlem myopathy, an autosomal dominant myopathy with contractures. *Nat Genet* 14:113-15.
- Jones FS, Jones PL (2000): The tenascin family of ECM glycoproteins: structure, function, and regulation during embryonic development and tissue remodeling. *Dev Dyn* 218:235-59.
- Jones JC, Hopkinson SB, Goldfinger LE (1998): Structure and assembly of hemidesmosomes. *Bioessays* 20:488-94.
- Jones PL, Jones FS, Zhou B, Rabinovitch M (1999): Induction of vascular smooth muscle cell tenascin-C gene expression by denatured type I collagen

- is dependent upon a $\beta 3$ integrin-mediated mitogen-activated protein kinase pathway and a 122-base pair promoter element. *J Cell Sci* 112 (Pt 4): 435-45.
- Kadler K (1994): Extracellular matrix. 1: Fibril-forming collagens. *Protein Profile* 1:519-638.
- Kadler K (1995): Extracellular matrix 1: Fibril-forming collagens. *Protein Profile* 2:491-619.
- Kashtan CE (1995): Clinical and molecular diagnosis of Alport syndrome. *Proc Assoc Am Physicians* 107:306-13.
- Kashtan CE (2000): Alport syndrome: Abnormalities of type IV collagen genes and proteins. *Ren Fail* 22:737-49.
- Khwaja A, Rodriguez-Viciano P, Wennstrom S, Warne PH, Downward J (1997): Matrix adhesion and Ras transformation both activate a phosphoinositide 3-OH kinase and protein kinase B/Akt cellular survival pathway. *EMBO J* 16:2783-93.
- Kleinman HK, McGarvey ML, Liotta LA, Robey PG, Tryggvason K, Martin GR (1982): Isolation and characterization of type IV procollagen, laminin, and heparan sulfate proteoglycan from the EHS sarcoma. *Biochemistry* 21: 6188-93.
- Kon A, Pulkkinen L, Hara M, et al. (1998): Laminin 5 genes and Herlitz junctional epidermolysis bullosa: novel mutations and polymorphisms in the LAMB3 and LAMC2 genes: Mutations in brief no. 190. *Online. Hum Mutat* 12:288.
- Kreidberg JA, Donovan MJ, Goldstein SL, et al. (1996): Alpha 3 beta 1 integrin has a crucial role in kidney and lung organogenesis. *Development* 122: 3537-47.
- Kresse H, Schonherr E (2001): Proteoglycans of the extracellular matrix and growth control. *J Cell Physiol* 189:266-74.
- Kuang W, Xu H, Vachon PH, et al. (1998): Merosin-deficient congenital muscular dystrophy: Partial genetic correction in two mouse models. *J Clin Invest* 102:844-52.
- Kuhn K (1995): Basement membrane (type IV) collagen. *Matrix Biol* 14:439-45.
- Kuivaniemi H, Tromp G, Prockop DJ (1997): Mutations in fibrillar collagens (types I, II, III, and XI), fibril-associated collagen (type IX), and network-forming collagen (type X) cause a spectrum of diseases of bone, cartilage, and blood vessels. *Hum Mutat* 9:300-15.
- Kuster JE, Guarnieri MH, Ault JG, Flaherty L, Swiatek PJ (1997): IAP insertion in the murine Lamb3 gene results in junctional epidermolysis bullosa. *Mamm Genome* 8:673-81.
- Lander AD, Stipp CS, Ivins JK (1996): The glypican family of heparan sulfate proteoglycans: Major cell-surface proteoglycans of the developing nervous system. *Perspect Dev Neurobiol* 3:347-58.
- Law DA, DeGuzman FR, Heiser P, Ministri-Madrid K, Killeen N, Phillips DR (1999): Integrin cytoplasmic tyrosine motif is required for outside-in α IIb β 3 signalling and platelet function. *Nature* 401:808-11.
- Le Gall M, Grall D, Chambard JC, Pouyssegur J, Van Obberghen-Schilling E (1998): An anchorage-dependent signal distinct from p42/44 MAP kinase activation is required for cell cycle progression. *Oncogene* 17:1271-7.

- Lemmink HH, Nillesen WN, Mochizuki T, et al. (1996): Benign familial hematuria due to mutation of the type IV collagen alpha4 gene. *J Clin Invest* 98: 1114-18.
- Lewis JM, Schwartz MA (1995): Mapping in vivo associations of cytoplasmic proteins with integrin beta 1 cytoplasmic domain mutants. *Mol Biol Cell* 6: 151-60.
- Li J, Yen C, Liaw D, et al. (1997): PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 275:1943-7.
- Li ML, Aggeler J, Farson DA, Hatier C, Hassell J, Bissell MJ (1987): Influence of a reconstituted basement membrane and its components on casein gene expression and secretion in mouse mammary epithelial cells. *Proc Natl Acad Sci USA* 84:136-40.
- Li Y, Lacerda DA, Warman ML, et al. (1995): A fibrillar collagen gene, *Col11a1*, is essential for skeletal morphogenesis. *Cell* 80:423-30.
- Lin CQ, Bissell MJ (1993): Multi-faceted regulation of cell differentiation by extracellular matrix. *Faseb J* 7:737-43.
- Lin TH, Chen Q, Howe A, Juliano RL (1997): Cell anchorage permits efficient signal transduction between *ras* and its downstream kinases. *J Biol Chem* 272: 8849-52.
- Littlewood Evans A, Muller U (2000): Stereocilia defects in the sensory hair cells of the inner ear in mice deficient in integrin alpha8beta1. *Nat Genet* 24:424-8.
- Liu F, Sells MA, Chernoff J (1998): Protein tyrosine phosphatase 1B negatively regulates integrin signaling. *Curr Biol* 8:173-6.
- Liu X, Wu H, Byrne M, Jeffrey J, Krane S, Jaenisch R (1995): A targeted mutation at the known collagenase cleavage site in mouse type I collagen impairs tissue remodeling. *J Cell Biol* 130:227-37.
- Liu X, Wu H, Byrne M, Krane S, Jaenisch R (1997): Type III collagen is crucial for collagen I fibrillogenesis and for normal cardiovascular development. *Proc Natl Acad Sci USA* 94:1852-6.
- Lohi J, Leivo I, Franssila K, Virtanen I (1997): Changes in the distribution of integrins and their basement membrane ligands during development of human thyroid follicular epithelium. *Histochem J* 29:337-45.
- Maehama T, Dixon JE (1998): The tumor suppressor, PTEN/MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *J Biol Chem* 273:13375-8.
- Mayer U, Kohfeldt E, Timpl R (1998): Structural and genetic analysis of laminin-nidogen interaction. *An NY Acad Sci* 857:130-42.
- Mayer U, Saher G, Fassler R, et al. (1997): Absence of integrin alpha 7 causes a novel form of muscular dystrophy. *Nat Genet* 17:318-23.
- McGuirt WT, Prasad SD, Griffith AJ, et al. (1999): Mutations in *COL11A2* cause non-syndromic hearing loss (DFNA13). *Nat Genet* 23:413-19.
- McHugh KP, Hodivala-Dilke K, Zheng MH, et al. (2000): Mice lacking beta3 integrins are osteosclerotic because of dysfunctional osteoclasts. *J Clin Invest* 105:433-40.
- McIntosh I, Abbott MH, Francomano CA (1995): Concentration of mutations causing Schmid metaphyseal chondrodysplasia in the C-terminal noncollagenous domain of type X collagen. *Hum Mutat* 5:121-5.
- Meek KM, Fullwood NJ (2001): Corneal and scleral collagens—A microscopist's perspective. *Micron* 32:261-72.

- Meisler MH, Griffith AJ, Warman M, Tiller G, Sprunger LK (1998): Gene symbol: COL11A1. Disease: Marshall syndrome. *Hum Genet* 102:498.
- Miosge N (2001): The ultrastructural composition of basement membranes in vivo. *Histol Histopathol* 16:1239-48.
- Miosge N, Klenczar C, Herken R, Willem M, Mayer U (1999): Organization of the myotendinous junction is dependent on the presence of $\alpha 7\beta 1$ integrin. *Lab Invest* 79:1591-9.
- Miranti CK, Brugge JS (2002): Sensing the environment: A historical perspective on integrin signal transduction. *Nat Cell Biol* 4:E83-90.
- Miyamoto S, Teramoto H, Coso OA, et al. (1995): Integrin function: Molecular hierarchies of cytoskeletal and signaling molecules. *J Cell Biol* 131:791-805.
- Miyamoto S, Teramoto H, Gutkind JS, Yamada KM (1996): Integrins can collaborate with growth factors for phosphorylation of receptor tyrosine kinases and MAP kinase activation: Roles of integrin aggregation and occupancy of receptors. *J Cell Biol* 135:1633-42.
- Moore SA, Saito F, Chen J, et al. (2002): Deletion of brain dystroglycan recapitulates aspects of congenital muscular dystrophy. *Nature* 418:422-5.
- Muller U, Wang D, Denda S, Meneses JJ, Pedersen RA, Reichardt LF (1997): Integrin $\alpha 8\beta 1$ is critically important for epithelial-mesenchymal interactions during kidney morphogenesis. *Cell* 88:603-13.
- Murgia C, Blaikie P, Kim N, Dans M, Petrie HT, Giancotti FG (1998): Cell cycle and adhesion defects in mice carrying a targeted deletion of the integrin $\beta 4$ cytoplasmic domain. *EMBO J* 17:3940-51.
- Murshed M, Smyth N, Miosge N, et al. (2000): The absence of nidogen 1 does not affect murine basement membrane formation. *Mol Cell Biol* 20:7007-12.
- Muschler J, Levy D, Boudreau R, Henry M, Campbell K, Bissell MJ (2002): A role for dystroglycan in epithelial polarization: loss of function in breast tumor cells. *Cancer Res* 62:7102-9.
- Muthuswamy SK, Li D, Lelievre S, Bissell MJ, Brugge JS (2001): ErbB2, but not ErbB1, reinitiates proliferation and induces luminal repopulation in epithelial acini. *Nat Cell Biol* 3:785-92.
- Myers MP, Pass I, Batty IH, et al. (1998): The lipid phosphatase activity of PTEN is critical for its tumor suppressor function. *Proc Natl Acad Sci USA* 95:13513-18.
- Myllyharju J, Kivirikko KI (2001): Collagens and collagen-related diseases. *An Med* 33:7-21.
- Nievers MG, Schaapveld RQ, Sonnenberg A (1999): Biology and function of hemidesmosomes. *Matrix Biol* 18:5-17.
- Noakes PG, Gautam M, Mudd J, Sanes JR, Merlie JP (1995): Aberrant differentiation of neuromuscular junctions in mice lacking α -laminin/laminin $\beta 2$. *Nature* 374:258-62.
- Oh ES, Gu H, Saxton TM, et al. (1999): Regulation of early events in integrin signaling by protein tyrosine phosphatase SHP-2. *Mol Cell Biol* 19:3205-15.
- Oldberg A, Antonsson P, Hedbom E, Heinegard D (1990): Structure and function of extracellular matrix proteoglycans. *Biochem Soc Trans* 18:789-92.
- Olsen BR (1995): New insights into the function of collagens from genetic analysis. *Curr Opin Cell Biol* 7:720-7.

- Paine-Saunders S, Viviano BL, Zupicich J, Skarnes WC, Saunders S (2000): Glypican-3 controls cellular responses to Bmp4 in limb patterning and skeletal development. *Dev Biol* 225:179-87.
- Park PW, Reizes O, Bernfield M (2000): Cell surface heparan sulfate proteoglycans: Selective regulators of ligand-receptor encounters. *J Biol Chem* 275:29923-6.
- Parsons JT, Schaller MD, Hildebrand J, Leu TH, Richardson A, Otey C (1994): Focal adhesion kinase: Structure and signalling. *J Cell Sci Suppl* 18:109-13.
- Patton BL, Cunningham JM, Thyboll J, et al. (2001): Properly formed but improperly localized synaptic specializations in the absence of laminin alpha4. *Nat Neurosci* 4:597-604.
- Perrimon N, Bernfield M (2001): Cellular functions of proteoglycans—An overview. *Semin Cell Dev Biol* 12:65-7.
- Petersen OW, Ronnov-Jessen L, Howlett AR, Bissell MJ (1992): Interaction with basement membrane serves to rapidly distinguish growth and differentiation pattern of normal and malignant human breast epithelial cells. *Proc Natl Acad Sci USA* 89:9064-8.
- Petitclerc E, Boutaud A, Prestayko A, et al. (2000): New functions for non-collagenous domains of human collagen type IV: Novel integrin ligands inhibiting angiogenesis and tumor growth in vivo. *J Biol Chem* 275:8051-61.
- Phillips CL, Pfeiffer BJ, Luger AM, Franklin CL (2002): Novel collagen glomerulopathy in a homotrimeric type I collagen mouse (oim). *Kidney Int* 62:383-91.
- Plopper G, Ingber DE (1993): Rapid induction and isolation of focal adhesion complexes. *Biochem Biophys Res Commun* 193:571-8.
- Plopper GE, McNamee HP, Dike LE, Bojanowski K, Ingber DE (1995): Convergence of integrin and growth factor receptor signaling pathways within the focal adhesion complex. *Mol Biol Cell* 6:1349-65.
- Pozzi A, Wary KK, Giancotti FG, Gardner HA (1998): Integrin alpha1beta1 mediates a unique collagen-dependent proliferation pathway in vivo. *J Cell Biol* 142:587-94.
- Pulkkinen L, Kim DU, Uitto J (1998): Epidermolysis bullosa with pyloric atresia: novel mutations in the beta4 integrin gene (ITGB4). *Am J Pathol* 152:157-66.
- Radisky D, Muschler J, Bissell MJ (2002): Order and disorder: The role of extracellular matrix in epithelial cancer. *Cancer Invest* 20:139-53.
- Raghavan S, Bauer C, Mundschauf G, Li Q, Fuchs E (2000): Conditional ablation of beta1 integrin in skin: Severe defects in epidermal proliferation, basement membrane formation, and hair follicle invagination. *J Cell Biol* 150:1149-60.
- Rapraeger AC (2000): Syndecan-regulated receptor signaling. *J Cell Biol* 149:995-8.
- Reizes O, Lincecum J, Wang Z, et al. (2001): Transgenic expression of syndecan-1 uncovers a physiological control of feeding behavior by syndecan-3. *Cell* 106:105-16.
- Ren XD, Kiousses WB, Schwartz MA (1999): Regulation of the small GTP-binding protein Rho by cell adhesion and the cytoskeleton. *EMBO J* 18:578-85.
- Ren XD, Kiousses WB, Sieg DJ, Otey CA, Schlaepfer DD, Schwartz MA (2000): Focal adhesion kinase suppresses Rho activity to promote focal adhesion turnover. *J Cell Sci* 113 (Pt 20):3673-8.
- Renshaw MW, Price LS, Schwartz MA (1999): Focal adhesion kinase mediates the integrin signaling requirement for growth factor activation of MAP kinase. *J Cell Biol* 147:611-8.

- Richards AJ, Martin S, Yates JR, et al. (2000): COL2A1 exon 2 mutations: Relevance to the Stickler and Wagner syndromes. *Br J Ophthalmol* 84:364-71.
- Ritvaniemi P, Korkko J, Bonaventure J, et al. (1995): Identification of COL2A1 gene mutations in patients with chondrodysplasias and familial osteoarthritis. *Arthritis Rheum* 38:999-1004.
- Rock MT, Brooks WH, Roszman TL (1997): Calcium-dependent signaling pathways in T cells: Potential role of calpain, protein tyrosine phosphatase 1b, and p130Cas in integrin-mediated signaling events. *J Biol Chem* 272:33377-83.
- Ronnov-Jessen L, Petersen OW, Bissell MJ (1996): Cellular changes involved in conversion of normal to malignant breast: Importance of the stromal reaction. *Physiol Rev* 76:69-125.
- Roovers K, Assoian RK (2000): Integrating the MAP kinase signal into the G₁ phase cell cycle machinery. *Bioessays* 22:818-26.
- Roovers K, Davey G, Zhu X, Bottazzi ME, Assoian RK (1999): Alpha5beta1 integrin controls cyclin D1 expression by sustaining mitogen-activated protein kinase activity in growth factor-treated cells. *Mol Biol Cell* 10:3197-204.
- Rossino P, Gavazzi I, Timpl R, et al. (1990): Nerve growth factor induces increased expression of a laminin-binding integrin in rat pheochromocytoma PC12 cells. *Expr Cell Res* 189:100-8.
- Ryan MC, Lee K, Miyashita Y, Carter WG (1999): Targeted disruption of the LAMA3 gene in mice reveals abnormalities in survival and late stage differentiation of epithelial cells. *J Cell Biol* 145:1309-23.
- Schaller MD, Hildebrand JD, Shannon JD, Fox JW, Vines RR, Parsons JT (1994): Autophosphorylation of the focal adhesion kinase, pp125FAK, directs SH2-dependent binding of pp60src. *Mol Cell Biol* 14:1680-8.
- Schaller MD, Otey CA, Hildebrand JD, Parsons JT (1995): Focal adhesion kinase and paxillin bind to peptides mimicking beta integrin cytoplasmic domains. *J Cell Biol* 130:1181-7.
- Schlaepfer DD, Hanks SK, Hunter T, van der Geer P (1994): Integrin-mediated signal transduction linked to Ras pathway by GRB2 binding to focal adhesion kinase. *Nature* 372:786-91.
- Schon MP, Schon M, Warren HB, Donohue JP, Parker CM (2000): Cutaneous inflammatory disorder in integrin alphaE (CD103)-deficient mice. *J Immunol* 165:6583-9.
- Schwartz MA (1997): Integrins, oncogenes, and anchorage independence. *J Cell Biol* 139:575-8.
- Schwartz MA, Assoian RK (2001): Integrins and cell proliferation: Regulation of cyclin-dependent kinases via cytoplasmic signaling pathways. *J Cell Sci* 114:2553-60.
- Schwartz MA, Assoian RK (2001): Integrins and cell proliferation: Regulation of cyclin-dependent kinases via cytoplasmic signaling pathways. *J Cell Sci* 114:2553-60.
- Schwartz MA, Ingber DE (1994): Integrating with integrins. *Mol Biol Cell* 5:389-93.
- Schwartz MA, Schaller MD, Ginsberg MH (1995): Integrins: Emerging paradigms of signal transduction. *An Rev Cell Dev Biol* 11:549-99.
- Schymeinsky J, Nedbal S, Miosge N, et al. (2002): Gene structure and functional analysis of the mouse nidogen-2 gene: Nidogen-2 is not essential for basement membrane formation in mice. *Mol Cell Biol* 22:6820-30.

- Sherr CJ (1994): G₁ phase progression: Cycling on cue. *Cell* 79:551-5.
- Simpson DG, Terracio L, Terracio M, Price RL, Turner DC, Borg TK (1994): Modulation of cardiac myocyte phenotype in vitro by the composition and orientation of the extracellular matrix. *J Cell Physiol* 161:89-105.
- Smith LT, Schwarze U, Goldstein J, Byers PH (1997): Mutations in the COL3A1 gene result in the Ehlers-Danlos syndrome type IV and alterations in the size and distribution of the major collagen fibrils of the dermis. *J Invest Dermatol* 108:241-7.
- Smyth N, Vatansever HS, Murray P, et al. (1999): Absence of basement membranes after targeting the LAMC1 gene results in embryonic lethality due to failure of endoderm differentiation. *J Cell Biol* 144:151-60.
- Snead MP, Yates JR, Williams R, Payne SJ, Pope FM, Scott JD (1996): Stickler syndrome type 2 and linkage to the COL11A1 gene. *Am NY Acad Sci* 785: 331-2.
- Spranger J, Menger H, Mundlos S, Winterpacht A, Zabel B (1994): Kniest dysplasia is caused by dominant collagen II (COL2A1) mutations: Parental somatic mosaicism manifesting as Stickler phenotype and mild spondyloepiphyseal dysplasia. *Pediatr Radiol* 24:431-5.
- Stambolic V, Suzuki A, de la Pompa JL, et al. (1998): Negative regulation of PKB/Akt-dependent cell survival by the tumor suppressor PTEN. *Cell* 95: 29-39.
- Stephens LE, Sutherland AE, Klimanskaya IV, et al. (1995): Deletion of beta 1 integrins in mice results in inner cell mass failure and peri-implantation lethality. *Genes Dev* 9:1883-95.
- Sternlicht MD, Bissell MJ, Werb Z (2000): The matrix metalloproteinase stromelysin-1 acts as a natural mammary tumor promoter. *Oncogene* 19: 1102-13.
- Sternlicht MD, Lochter A, Sympon CJ, et al. (1999): The stromal proteinase MMP3/stromelysin-1 promotes mammary carcinogenesis. *Cell* 98:137-46.
- Streuli C (1999): Extracellular matrix remodelling and cellular differentiation. *Curr Opin Cell Biol* 11:634-40.
- Su J, Muranjan M, Sap J (1999): Receptor protein tyrosine phosphatase alpha activates Src-family kinases and controls integrin-mediated responses in fibroblasts. *Curr Biol* 9:505-11.
- Takuwa N, Fukui Y, Takuwa Y (1999): Cyclin D1 expression mediated by phosphatidylinositol 3-kinase through mTOR-p70(S6K)-independent signaling in growth factor-stimulated NIH 3T3 fibroblasts. *Mol Cell Biol* 19:1346-58.
- Talhouk RS, Bissell MJ, Werb Z (1992): Coordinated expression of extracellular matrix-degrading proteinases and their inhibitors regulates mammary epithelial function during involution. *J Cell Biol* 118:1271-82.
- Tamura M, Gu J, Matsumoto K, Aota S, Parsons R, Yamada KM (1998): Inhibition of cell migration, spreading, and focal adhesions by tumor suppressor PTEN. *Science* 280:1614-17.
- Tiller GE, Polumbo PA, Weis MA, et al. (1995): Dominant mutations in the type II collagen gene, COL2A1, produce spondyloepimetaphyseal dysplasia, Strudwick type. *Nat Genet* 11:87-9.
- Timpl R, Fujiwara S, Dziadek M, Aumailley M, Weber S, Engel J (1984): Laminin, proteoglycan, nidogen and collagen IV: Structural models and molecular interactions. *Ciba Found Symp* 108:25-43.

- Toole BP (2001): Hyaluronan in morphogenesis. *Semin Cell Dev Biol* 12:79-87.
- Toole BP (2002): Hyaluronan promotes the malignant phenotype. *Glycobiology* 12:37R-42R.
- Toole BP, Wight TN, Tammi MI (2002): Hyaluronan-cell interactions in cancer and vascular disease. *J Biol Chem* 277:4593-6.
- Trummer T, Brenner R, Just W, Vogel W, Kennerknecht I (2001): Recurrent mutations in the COL1A2 gene in patients with osteogenesis imperfecta. *Clin Genet* 59:338-43.
- Tryggvason K (1995): Molecular properties and diseases of collagens. *Kidney Int Suppl* 49:S24-8.
- Tsai LH (1998): Stuck on the ECM. *Trends Cell Biol* 8:292-5.
- Turley EA, Noble PW, Bourguignon LY (2002): Signaling properties of hyaluronan receptors. *J Biol Chem* 277:4589-92.
- Uitto J, Pulkkinen L (1996): Molecular complexity of the cutaneous basement membrane zone. *Mol Biol Rep* 23:35-46.
- Vachon PH, Loechel F, Xu H, Wewer UM, Engvall E (1996): Merosin and laminin in myogenesis; specific requirement for merosin in myotube stability and survival. *J Cell Biol* 134:1483-97.
- van Keulen CJ, van de Akker E, Pals G, Raurwerda JA (1999): The role of type III collagen in the development of familial abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 18:65-70.
- Vidal F, Baudoin C, Miquel C, et al. (1995): Cloning of the laminin alpha 3 chain gene (LAMA3) and identification of a homozygous deletion in a patient with Herlitz junctional epidermolysis bullosa. *Genomics* 30:273-80.
- Virtanen I, Laitinen L, Korhonen M (1995): Differential expression of laminin polypeptides in developing and adult human kidney. *J Histochem Cytochem* 43:621-8.
- Vuori K, Hirai H, Aizawa S, Ruoslahti E (1996): Introduction of p130cas signaling complex formation upon integrin-mediated cell adhesion: A role for Src family kinases. *Mol Cell Biol* 16:2606-13.
- Wagner N, Lohler J, Kunkel EJ, et al. (1996): Critical role for beta7 integrins in formation of the gut-associated lymphoid tissue. *Nature* 382:366-70.
- Wai AW, Ng LJ, Watanabe H, Yamada Y, Tam PP, Cheah KS (1998): Disrupted expression of matrix genes in the growth plate of the mouse cartilage matrix deficiency (cmd) mutant. *Dev Genet* 22:349-58.
- Wang F, Weaver VM, Petersen OW, et al. (1998): Reciprocal interactions between beta1-integrin and epidermal growth factor receptor in three-dimensional basement membrane breast cultures: A different perspective in epithelial biology. *Proc Natl Acad Sci USA* 95:14821-6.
- Ward LM, Lalic L, Roughley PJ, Glorieux FH (2001): Thirty-three novel COL1A1 and COL1A2 mutations in patients with osteogenesis imperfecta types I-IV. *Hum Mutat* 17:434.
- Wary KK, Dans M, Mariotti A, Giancotti FG (1999): Biochemical analysis of integrin-mediated Shc signaling. *Methods Mol Biol* 129:35-49.
- Wary KK, Mainiero F, Isakoff SJ, Marcantonio EE, Giancotti FG (1996): The adaptor protein Shc couples a class of integrins to the control of cell cycle progression. *Cell* 87:733-43.
- Wasylyk C, Gutman A, Nicholson R, Wasylyk B (1991): The c-Ets oncoprotein activates the stromelysin promoter through the same elements as several non-nuclear oncoproteins. *EMBO J* 10:1127-34.

- Watt FM (2002): Role of integrins in regulating epidermal adhesion, growth and differentiation. *EMBO J* 21:3919-26.
- Weaver VM, Howlett AR, Langton-Webster B, Petersen OW, Bissell MJ (1995): The development of a functionally relevant cell culture model of progressive human breast cancer. *Semin Cancer Biol* 6:175-84.
- Weaver VM, Petersen OW, Wang F, et al. (1997): Reversion of the malignant phenotype of human breast cells in three-dimensional culture and in vivo by integrin blocking antibodies. *J Cell Biol* 137:231-45.
- Weber JD, Raben DM, Phillips PJ, Baldassare JJ (1997): Sustained activation of extracellular-signal-regulated kinase 1 (ERK1) is required for the continued expression of cyclin D1 in G₁ phase. *Biochem J* 326 (Pt 1):61-8.
- Wei Y, Yang X, Liu Q, Wilkins JA, Chapman HA (1999): A role for caveolin and the urokinase receptor in integrin-mediated adhesion and signaling. *J Cell Biol* 144:1285-94.
- Weinberg RA (1995): The retinoblastoma protein and cell cycle control. *Cell* 81:323-30.
- Werb Z, Chin JR (1998): Extracellular matrix remodeling during morphogenesis. *An NY Acad Sci* 857:110-18.
- Wewer UM, Engvall E (1996): Merosin/laminin-2 and muscular dystrophy. *Neuromuscul Disord* 6:409-18.
- Williamson RA, Henry MD, Daniels KJ, et al. (1997): Dystroglycan is essential for early embryonic development: Disruption of Reichert's membrane in Dag1-null mice. *Hum Mol Genet* 6:831-41.
- Wiseman BS, Werb Z (2002): Stromal effects on mammary gland development and breast cancer. *Science* 296:1046-9.
- Woods D, Cherwinski H, Venetsanakos E, et al. (2001): Induction of beta3-integrin gene expression by sustained activation of the Ras-regulated Raf-MEK-extracellular signal-regulated kinase signaling pathway. *Mol Cell Biol* 21:3192-205.
- Xu H, Wu XR, Wewer UM, Engvall E (1994): Murine muscular dystrophy caused by a mutation in the laminin alpha 2 (Lama2) gene. *Nat Genet* 8:297-302.
- Yamakita Y, Totsukawa G, Yamashiro S, et al. (1999): Dissociation of FAK/p130(CAS)/c-Src complex during mitosis: Role of mitosis-specific serine phosphorylation of FAK. *J Cell Biol* 144:315-24.
- Yanagishita M (1993): Function of proteoglycans in the extracellular matrix. *Acta Pathol Jpn* 43:283-93.
- Yang JT, Rayburn H, Hynes RO (1993): Embryonic mesodermal defects in alpha 5 integrin-deficient mice. *Development* 119:1093-105.
- Yang JT, Rayburn H, Hynes RO (1995): Cell adhesion events mediated by alpha 4 integrins are essential in placental and cardiac development. *Development* 121:549-60.
- Yurchenco PD, Cheng YS, Colognato H (1992): Laminin forms an independent network in basement membranes. *J Cell Biol* 117:1119-33.
- Zhu X, Ohtsubo M, Bohmer RM, Roberts JM, Assoian RK (1996): Adhesion-dependent cell cycle progression linked to the expression of cyclin D1, activation of cyclin E-cdk2, and phosphorylation of the retinoblastoma protein. *J Cell Biol* 133:391-403.
- Zimmermann P, David G (1999): The syndecans, tuners of transmembrane signaling. *Faseb J* 13 Suppl:S91-S100.